
Energy Frontier

Conveners: R. Brock and M. E. Peskin

[Tags, e.g. (Q#1), refer to items on the Big Question list.]

1.1 Energy Frontier

The first run of the Large Hadron Collider (LHC) at CERN has closed a nearly half-century old chapter in the story of elementary particle physics. We have discovered a most unusual, new particle with properties very similar to those expected of the Higgs boson. The appearance of this particle—and further confirmation of its identity—closes one era and opens another. On one hand, the Standard Model of particle physics is complete. We know all of the particles in this model and how they interact with one another and we have at least a basic idea of their properties. On the other hand, we also know that the Standard Model is *incomplete* in important ways. It challenges us to uncover the physics behind its apparently ad hoc structure. We are certain that a host of observed anomalous phenomena and set of confusing conceptual questions have explanations that require new physics outside the Standard Model.

The LHC and the CMS, ATLAS, and LHCb detectors have brought to bear impressive capabilities for exploring for the answers to these new questions. The LHC accelerator is expected to dramatically increase its ability to deliver beams in the period between now and 2030, increasing its energy by almost a factor of two and its integrated luminosity by a factor of 100. In concert, the detectors will improve their ability to collect enormous data sets and to discriminate the properties of events with increasing precision. Further, around the world, other new accelerators are being considered that will give us additional power in understanding the heaviest particles of the Standard Model and exploring for new ones. In this report, and in the detailed working group reports, we trace out the programs of these accelerators and present their most important goals.

Our successful theory of weak interactions is based on the idea of an underlying symmetry that is spontaneously broken. The symmetry of the theory of weak interactions dictates the couplings of the quarks and leptons to the W and Z bosons in a structure that has been confirmed by high-precision experiments. However, this symmetry forbids the quarks, leptons, and vector bosons from having mass. The inclusion of mass within the Standard Model requires a condensate of fields that fills the universe. The Higgs field—the field of which the Higgs particle is the quantum—is the simplest realization of this idea. The problem with this model is that it is too simple. These ideas raise many questions, and the model does not answer them.

The primary motivation for performing experiments at high energy accelerators comes from the demand that we unravel the mysteries associated with the Higgs field (Q#1). We have no understanding of the mechanism behind the Higgs field condensation. We have no understanding of the origin of the couplings of quarks and leptons to the condensate. We have no idea how to compute the couplings of the Higgs to quarks and leptons nor the potential energy that leads to its condensation. Finally, all of the Standard Model particles come in groups, yet the Higgs boson stands alone with properties that are unlike those of any other particle. This would be a puzzle in any case, but it also emphasizes that important pieces of the story are

missing. If these properties of the Higgs field and its quantum have physics explanations, then there must be new particles and forces, and their masses cannot be far above the mass scale of the Higgs boson itself.

These puzzles imply that new particles with masses of the order of 1 TeV which resolve these questions will be found—and will be accessible to existing and planned accelerators. (Q#9). The discovery of the Higgs particle and acceptance of the challenges that it poses obligate us to carry out a three-prong program of research in colliding beam experiments:

1. First, we must search for the direct production of these particles at high energy.
2. Second, we must search for the imprint of these particles on the couplings the W and Z bosons and the top quark.
3. Finally, we must study the Higgs boson itself in as much detail as possible, searching for signs of a larger Higgs sector structure and the influence of new heavy particles.

The Energy Frontier study pointed to all three of these approaches as important motivations for further experiments at colliders. The results of the study confirmed that the existing LHC detectors and their planned upgrades, together with proposed precision lepton collider experiments, will be nimble enough and sensitive enough to carry this three-fold campaign forward into the next two decades.

The Energy Frontier study was organized into six working groups, each associated with a physics topic. Each working group was asked to evaluate the future program for its subject both from a high-level perspective and from the viewpoint of supplying motivation for experiments at a range of proposed accelerators. In the remainder of this section, we will present the conclusions of these reports, first by physics topic, then by facility.

We begin with the Higgs boson. The bosonic resonance at 125 GeV was discovered at the LHC only one year ago. Its properties are now known to about the 30% level and, up to this point, its properties are consistent with those of the Higgs boson of the minimal Standard Model. The couplings of this boson roughly scale with mass. The specific form of the coupling to the Z boson indicates that the particle has spin-parity 0^+ and that the corresponding field has a nonzero vacuum expectation value.

However, we cannot be complacent about the identity and role of this particle (Q#1). On one hand, the idea that a single scalar field is solely responsible for the generation of all particle masses is just a guess and needs explicit verification. On the other hand, models with additional Higgs bosons and related new particles, and models in which the Higgs boson is composite, are hardly tested. Deviations from the minimal Higgs boson properties due to new particles with mass M are suppressed by a factor $(m_h/M)^2$, so to the extent that the LHC has set lower limits on the masses of new particles at many hundreds of GeV, we would not yet expect to see the modifications to the Higgs properties caused by those particles.

An experimental program to probe the Higgs boson contains several elements. The first is to search for deviations from the minimal Standard Model expectation that the Higgs boson couples to each particle species according to its mass. Such deviations are expected in almost all models of new physics, with a characteristic pattern for each model. The High-Luminosity LHC is expected to measure these couplings with accuracies of several percent, varying from coupling to coupling. Lepton collider experiments have the potential to push these accuracies to the sub-percent level, allowing discovery of percent-level deviations. Such a program of precision measurement of Higgs couplings requires a parallel concerted effort in precision theory. It also requires improvement of our knowledge of crucial input parameters such as α_s and m_b , which can be provided by lattice gauge theory computations. Collider experiments can also probe the nonlinear Higgs field self-coupling to the 10-20% level, thereby testing the critically important question of the shape of the Higgs potential.

Future experiments on the Higgs particle should also improve our knowledge of its mass and quantum numbers. The spin of the observed resonance should already be clear from LHC data in this decade. A more subtle question is whether this particle contains a small admixture of a CP-odd state, signaling CP violation in the Higgs sector (Q#4) and confirmation of at least one additional Higgs-like particle. We discuss probes for this effect at various colliders.

Finally, it is important to search directly for additional Higgs bosons (Q#9). The LHC can probe to masses of 1 TeV with model-dependent limits. Lepton colliders can probe more model-independently to masses close to $E_{CM}/2$.

QCD is well established as the correct theory of the strong interactions. Nevertheless, advances in QCD are needed to achieve the goals of future experiments, especially at hadron colliders. For experiments at hadron colliders, improved knowledge of the parton distribution functions is needed. This can be achieved with data expected from the LHC on the rapidity distribution of W , Z , and top quark production. In addition, precision cross section computations, to the NNLO level, are needed for many 2- and 3-particle production processes, especially those involving the Higgs boson. This will require advances in the theoretical art of QCD computation. Finally, it is important to push the error on the value of α_s below the 0.5% level. Lattice gauge theory seems to be the best avenue for achieving this.

The study of W and Z bosons has two aspects, first, the extension of the program of precision electroweak measurements, and, second, the search for perturbations of the three- and four-vector couplings that would signal the influence of new physics.

The minimal Standard Model makes precise predictions for the well-studied precision observables m_W and $\sin^2 \theta_w$ (Q#8). At the moment, the observed values are within 2σ of the predictions; the deviations are consistent with the effects of new particles in a range of new physics models. Higher accuracy in this program is clearly needed (Q#9). Future experiments will sharpen our knowledge of these quantities and perhaps expose inconsistency with the Standard Model. The LHC, especially in its high-luminosity phase, has the potential to reduce the error on the W mass to ± 5 MeV/ c^2 . This requires a factor 7 decrease in the current error due to parton distribution functions and is a challenge to QCD researchers. Lepton colliders can make further improvements, to an error of ± 2.5 MeV/ c^2 with a dedicated measurement of the WW threshold. A linear collider with beam polarization running at the Z resonance to produce 10^9 Z bosons (Giga- Z) is expected to reduce the error on $\sin^2 \theta_w$ by a factor 10. Finally, a circular e^+e^- collider operating at with synchrotron radiation luminosity in a 100 km tunnel can potentially push both errors down by another factor 4. All of these precision measurements and their interpretation push on the inflexible correlations among the Standard Model particles and their respective forces. Such precision measurements in precision electroweak observables could be discoveries should the tight constraints within the Standard Model begin to unravel.

The second theme of W and Z studies is the search for anomalous nonlinear couplings of the vector bosons (Q#8). Collider experiments at higher energy are sensitive to the triple gauge boson couplings and for the first time, to four-gauge-boson interactions, that is, direct measurement of vector boson scattering. Lepton collider experiments have the potential to push current uncertainties down by an order of magnitude, into the region in which new physics effects are predicted in models in which the Higgs boson is composite. Both hadron and lepton colliders can access vector boson scattering, but the total CM energy available to a two-vector-boson system is the crucial factor. The high-luminosity LHC is sensitive to vector boson or Higgs resonances with masses well above 1 TeV.

The top quark was discovered at the Fermilab Tevatron and studied there with samples of a few tens of thousands of $t\bar{t}$ pairs. At the LHC, we will study top quarks in samples of billions. At future lepton colliders, we will use the electroweak couplings of top quarks as a production mode and probe these with polarization observables. Both methods will transform our knowledge of this quark, whose properties are intimately

connected to the mysteries of flavor and mass generation. To this day, we are surprised at the extremely high mass of this presumably fundamental particle and its proximity to the value of the Higgs vacuum expectation value.

Although a puzzle in and of itself, the top quark mass is an important input parameter for particle physics (Q#2). The strongest demands on precision in the top quark mass come from the precision electroweak program, where interpretation of a 5 MeV/c² error in m_W requires a 600 MeV/c² error on m_t . This mass must be a theoretically well-defined quantity, convertible to a short-distance parameter such as the \overline{MS} mass. There are strategies applicable at the LHC which allow the measurement of a well-defined top quark mass to this 500 MeV/c² accuracy. At lepton colliders, measurement of the cross section at the top quark pair production threshold gives the \overline{MS} mass to 100 MeV/c², as required for the more accurate precision electroweak program available at these machines.

Top quark couplings will be studied at high accuracy both at hadron and at lepton colliders. New physics from top quark or Higgs compositeness can create few-percent corrections to the gluon, photon, and, especially, Z boson couplings. These effects can be observed as corrections to the pair-production cross sections from the predictions of the Standard Model. The top-quark coupling capabilities of a lepton collider is especially strong, with accuracies possible at the sub-percent level (Q#1). The billions of top quarks produced at the high-luminosity LHC allow very deep studies of rare flavor-changing top decays, to a level that complements searches at low energy for flavor-changing quark decays (Q#2).

Models of the Higgs potential and its symmetry breaking typically require new particles that are partners, in some way, of the top quark (Q#1). The LHC, especially in its high-luminosity stage, will have the capability for deep searches for supersymmetric partners of the top quark, heavy vector-like top quarks that appear in models with Higgs compositeness, and heavy resonances that decay to $t\bar{t}$, which appear in models with new space dimensions (Q#9).

High energy colliders can search more generally for new particles with a very broad range of properties (Q#9). These particles are required — and required to have masses near the 1 TeV scale — in models of electroweak symmetry breaking. Others among our questions also call for new particles accessible to high energy colliders. A large class of models of dark matter put the dark matter particle as the lightest particle of a TeV mass spectroscopy (Q#5). Grand unification requires new particles near the TeV scale, including partners of known particles and perhaps also new vector bosons associated with enhanced gauge symmetry (Q#8). CP violation in the Higgs boson sector is required in models that generate the matter-antimatter asymmetry at the electroweak phase transition (Q#4). More generally, new particles can bring new sources of flavor and CP violation that might be reflected in the discovery of new flavor-changing reactions at low energy (Q#2).

The LHC has already, in only its first run, increased the reach and power of searches for new particles over a broad scope. We expect that this power will increase dramatically in the next decade, as the LHC experiments acquire 300 fb⁻¹ of data at 14 TeV. This extension probes deeply into the region expected for new particles masses in all classes of models of electroweak symmetry breaking — and so, any plan for high energy physics in the longer term must include the possibility of new particles discovery in this period and exploitation of this discovery at the facilities that will follow. The high luminosity stage of the LHC, up to 3000 fb⁻¹, will provide a further very significant extension of the search region. This extension is particularly powerful for states produced through electroweak interactions, for which a factor of 2 increase in the mass reach is available in some case. Lepton colliders would bring new and complementary capabilities, in particular, the ability to carry out model-independent searches for states such as dark matter candidate particles whose signatures are especially difficult at hadron colliders (Q#5).

The physics opportunities described above are reflected as physics motivations for current and future high energy colliders. Our study considered a wide range of proposed machines; the full report from the Energy Frontier presents the cases for these machines in some detail. Here we have room for only the most important points.

We first point out the opportunity provided by the 14 TeV run of the LHC schedule for the next decade. This will provide robust searches for new particles over a broad front, with great promise of the discovery of the TeV particle spectrum motivated at the beginning of this section.

We find the case for the high-luminosity stage of the LHC compelling. This plan to deliver 3000 fb^{-1} , has been listed by the European Strategy Study as the highest priority accelerator project in Europe for the 2020's. We find that it will provide a significant additional step in the search for new particles, and that it will provide other important capabilities. The most important of these is the beginning of the era of precision Higgs boson measurements, to few-percent precision. It should give the first observation of the Higgs boson self-coupling. It will provide a program of precision measurement in the Standard Model that will dramatically tighten our knowledge of the W boson and the top quark, giving sensitivity in these dimensions to a variety of new physics models. While increasing the center of mass energy is obviously a significant leverage to higher mass states, we find that the factor of 10 integrated luminosity still increases the scale reach by 30% or more in many cases.

We considered the scientific case for the International Linear Collider. This next-stage lepton collider has recently completed its Technical Design Report and is judged at Snowmass to be ready for construction. This facility is named as the highest priority by the Japanese high energy physics community. We find the case for this machine to be very strong. It will reach sub-percent accuracy in the study of the Higgs boson, allowing discovery of percent-level effects in the Higgs couplings predicted in new physics models. It will measure the Higgs width in a model-independent way and will give the capability to observe Higgs coupling to dark matter and other invisible and exotic modes of Higgs decay. It will extend our knowledge of the top quark and the W and Z well below the percent level, setting up a confrontation with models that include Higgs boson and top quark composite structure.

Many other accelerator facilities have been proposed to the Snowmass study for construction over longer time scales. These include higher energy linear colliders, circular e^+e^- colliders, muon colliders, and photon colliders. We include a detailed discussion of the physics motivations for these facilities in our full report. There was particular interest in a proton collider of energy 100 TeV, which would come close to the capability of covering the full model space for models of "natural" electroweak symmetry breaking and WIMP dark matter. Our study developed materials and resources to begin a more complete study of physics at such a high energy collider.

Previous surveys of the prospects for high energy accelerator experiments have spoken in terms of reducing the space of parameters—couplings, mixings, masses—as if that were the goal. Now more than ever, the momentum points not toward exclusion, but toward the discovery of new states. Many possible directions are open and must be pursued.

The Higgs boson discovery changes everything. The study of this unusual particle to high precision, together with high precision studies of the W , Z , and top quark and searches for new states provide us with complementary routes into the mysterious physics of symmetry breaking. The current LHC detectors and their planned upgrades are well suited to carry on this program. Future accelerators will bring new capabilities to pursue it further.

High energy colliders provide manifest opportunities to discover new fundamental interactions of broad consequence. We believe that the US must be a part of these programs of discovery.