
Energy Frontier

Conveners: R. Brock and M. E. Peskin

K. Agashe, M. Artuso, J. Campbell, S. Dawson, R. Erbacher, C. Gerber, Y. Gershtein, A. Gribsan, K. Hatakeyama, J. Huston, A. Kotwal, H. Logan, M. Luty, K. Melnikov, M. Narain, M. Papucci, F. Petriello, S. Prell, J. Qian, R. Schwienhorst, C. Tully, R. Van Kooten, D. Wackerroth, L. Wang, D. Whiteson

October 3, 2013

1.1 Introduction

The original goal of elementary particle physics was to understand the nature of the subnuclear strong, electromagnetic, and weak forces. In the late 1960's and early 1970's, specific models for these forces were proposed in the form of Yang-Mills gauge theories, giving a beautiful explanation of all three interactions from a unified point of view. Together, these theories became known as “the Standard Model.” Today, we have a great deal of confidence that describing fundamental forces using the Gauge Principle is correct. Through precision experiments involving W and Z bosons carried out over the past twenty-five years, we have tested the Standard Model stringently, and the theory has passed every test. The most recent such experiments included the search for the Higgs boson, required in the Standard Model to generate quark, lepton, and vector boson masses. A year ago, the ATLAS and CMS experiments at the Large Hadron Collider discovered a candidate for this particle which, at the current level of our measurements, has all of the properties predicted in the Standard Model.

This is an historic level of success for theory and experiment: this economical model predicted the existence of fundamental fields, their dynamics, a scalar field responsible for the breaking of a gauge symmetry, and interactions among the particles with precision unmatched in the history of science. It all seems to have come true with remarkable accuracy. And yet, we find the result still unsatisfying. It is typically true in science that revolutionary changes in our understanding lead to a new set of vexing questions. The success of the Standard Model is no different. Though we have answered many questions about the structure of elementary particles, we have a new set of questions about the structure of the Standard Model itself. The discovery of the Higgs boson sharpens these issues and makes them even more mysterious.

There are many phenomena in nature that obviously lie outside of the Standard Model.

- We now know that 85% of the matter in the universe is *dark matter*—neutral, weakly interacting matter composed of one or more species not contained in the Standard Model.
- The cosmic excess of baryons over antibaryons is not explained by the Standard Model. Even though the Standard Model contains all of the necessary ingredients to generate baryon number in the early universe, including baryon number violation, CP violation, and a phase transition in cosmic history, the amount of baryon asymmetry generated is too small by ten orders of magnitude.

- 34 • The quantum numbers of the quarks and leptons under the Standard Model gauge symmetry $SU(3) \times$
35 $SU(2) \times U(1)$ strongly suggests that these symmetry groups are unified into a larger *grand unification*
36 group $SU(5)$ or $SO(10)$, our precision knowledge of the strengths of the gauge couplings is inconsistent
37 with this hypothesis.
- 38 • The Standard Model cannot account for neutrino masses without the addition of some new particles.
- 39 • Further, the pattern of weak interaction mixing among neutrinos is completely different from that
40 observed for quarks.
- 41 • The Standard Model does not include the force of gravity or the small but nonzero energy in empty
42 space that gives rise to *dark energy*.

43 The discovery of the Higgs boson has changed our viewpoint in how we address these questions, for three
44 reasons.

45
46 **First, the Higgs boson completes the particle spectrum of the Standard Model.** We now know all of the Standard
47 Model's ingredients and have at least a basic knowledge of their properties. It is clear now exactly what the
48 model does *not* explain. We have entered a new era in which the verification of the Standard Model takes
49 second place to a search for new, unknown forces and interactions.

50 **Second, one of the key mysteries concerns the Higgs boson itself.** The Higgs boson was predicted as a direct
51 consequence of the simplest model of the generation of mass for quarks, leptons, and gauge bosons. For
52 a long time, many particle physicists have expressed discomfort with this model. Now the prediction has
53 become a reality. We have to grapple with it and understand why nature chooses a particle with these
54 properties to do its work.

55 **Third, the Higgs boson itself gives us a new experimental approach.** As soon as we step outside the Standard
56 Model, the properties of the Higgs boson are hardly constrained by theory. It is compelling to tug on this
57 particle until the Standard Model breaks. We need to apply to the Higgs boson the same scrutiny that we
58 have applied in previous decades to hadron structure, heavy quark system, the W and Z bosons, and top
59 quark. Each study was done at the Energy Frontier machines of its day. This fruitful experimental approach
60 has acquired a new, promising target.

61 For exploration of the unknown regions outside the Standard Model, we are encouraged that very powerful,
62 experimental tools will be put into play. In the next ten years, the LHC at CERN is expected to almost
63 double its energy and to increase the size of its event sample by a factor of twenty. This new capability
64 will put to the test many models that predict new physics beyond the Standard Model and address the
65 unexplained phenomena listed earlier in this section. In the decade after that, the LHC should increase its
66 data set by a further factor of ten. Lepton colliders and higher energy hadron colliders are now receiving
67 serious consideration for construction. The mysteries associated with the Higgs boson call for new particles
68 and forces at the TeV energy scale or the attometer distance scale. Now we have before us capabilities for
69 a thorough exploration of this region of masses and distances. This is a compelling program; the purpose of
70 this report is to describe it in detail.

71 The structure of this summary report is as follows: In Section 2, we present the arguments for new
72 fundamental interactions at the TeV energy scale and the experimental program at colliders that these
73 arguments motivate. In Sections 3–8, we review in a more specific way the physics issues of collider
74 experiments at the TeV energy scale. We consider in turn the prospects for exploration of new physics
75 through studies of the Higgs boson, the W and Z bosons, Quantum Chromodynamics (QCD), the top
76 quark, and searches for and study of new particles. We present the questions that need to be answered

77 and the methodologies to attack these questions. In Section 9, we present the capabilities of current and
78 proposed colliders in relation to these physics goals. Section 10 gives our conclusions.

79 1.2 Importance of the TeV Scale

80 We have listed a number of motivations for new fundamental interactions beyond the Standard Model (SM).
81 Where will we find them?

82 Explanations for dark matter, baryogenesis, higher unification, and dark energy span a bewildering range
83 of mass and distance scale. However, many of the questions we have listed in the previous section relate
84 specifically to the energy scale of hundreds to thousands of GeV that we are exploring today at the Large
85 Hadron Collider. We consider it imperative to understand particles at forces at this “TeV scale” thoroughly,
86 using all of the tools at our disposal. In this section, we will discuss the importance of this regime of energies
87 and short distances.

88 There is a sharp boundary at which our well-founded knowledge of the fundamental elementary particle
89 interactions runs out. This is related to two different faces that the SM presents, which stand on very
90 different theoretical foundations. On one side are the Yang-Mills gauge interactions, on the other side, the
91 interactions of the Higgs field. The Yang-Mills interactions of quarks, leptons, and vector bosons are tightly
92 determined by their quantum numbers and the strength of the coupling constants of the $SU(3) \times SU(2) \times U(1)$
93 vector bosons. Precision tests of the SM confirm the structure of these interactions to impressive accuracy [1].
94 There is little doubt that, here, the SM is a correct representation of nature.

95 On the other hand, the interactions of the SM fermions with the Higgs field, and the dynamics of the
96 Higgs field itself, are essentially unconstrained and conceptually even cumbersome. The SM Lagrangian is
97 constructed by writing down the most general terms allowed by gauge symmetry and renormalizability. The
98 resulting potential term contains much of what is perplexing about the SM:

$$V = \mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2 + \sum_{f, f'} [G_{\Phi f f'} \bar{f}_L f'_R \Phi + h.c.] . \quad (1.1)$$

99 Here Φ is the spin 0 doublet field (doublet) and λ , $G_{\Phi f f'}$, and μ are parameters. The first two terms give
100 the potential energy of the Φ field. When $\mu^2 < 0$, the neutral $I_3 = -1/2$ component of Φ is singled out
101 and shifted by $\Phi^0 \rightarrow v + h$ where $v = \sqrt{-\mu^2/\lambda}$ is the vacuum expectation value. At the same time, we
102 diagonalize the matrix $G_{\Phi f f'}$. The last term in brackets then becomes

$$V(ff) = \sum_f \left[\frac{g_{Hff} v}{\sqrt{2}} \bar{f}_L f_R + \frac{g_{Hff}}{\sqrt{2}} \bar{f}_L f_R h + h.c. \right] . \quad (1.2)$$

103 Every term in (1.1) and (1.2) is of critical importance and each presents special challenges to interpretation
104 and measurement. The first term in (1.2) is the new mass term for fermions, $m_f = g_{Hff} v / \sqrt{2}$. The pattern
105 of the fermion masses is totally unconstrained and proportional to the coupling constant g_{Hff} . The Yukawa
106 fermion-Higgs couplings (the second term of (1.2) and the angles from the diagonalization) contain the origin
107 of mass and mixings among quarks and leptons and CP violation in the weak interactions. Many parameters
108 in this sector are well measured, but there is no theory that explains their origin and structure.

109 As to the couplings of the Higgs boson itself (the first two terms of 1.1), the picture given in the Standard
110 Model is just a guess. There may be additional Higgs bosons and additional particles of other types forming
111 a larger “Higgs sector,” we have almost no information about these particles except that their masses are
112 probably larger than the mass of the known Higgs boson at $\mu = 125 \text{ GeV}/c^2$.

1.2.1 The mystery of Higgs field symmetry breaking

The questions raised at the end of the previous section are brought to a focus by a single underlying question: In order for any SM quark, lepton, or vector boson to obtain mass, the Φ field must condense and the resultant Higgs field fills the universe. Why does this happen?

The SM itself provides no help with this question. It states only that symmetry breaking occurs if $\mu^2 < 0$, which, as a physics explanation, is completely empty. Potentials of the form of (1.1) appear in many condensed matter systems, including superconductors, magnets, and binary alloys. In those systems, it is possible to compute the parameters of the potential from the underlying details of atomic structure and explain why $\mu^2 < 0$. For the SM, if there is an underlying dynamics, its form is unknown. Attempts to compute μ^2 within the SM, even to determine its sign, give disastrous results. The answers for μ^2 depend quadratically on the values of large, unknown mass scales, with competing contributions of opposite sign.

Models are known in which μ^2 can be computed. However, they are not simple extensions of the SM. The barrier to such a model is that the quadratic dependence on unknown scale parameters at very high energy must be removed. However, this dependence is a generic property of models with fundamental scalar fields, associated with the fact that the radiative corrections to the scalar field mass are quadratically divergent. Cures for this problem require that the Higgs particle is non-generic in some important way: Either it is a composite particle or it is related by a symmetry to a fermion or a vector boson. Symmetries of these types can be included consistently only by profound extension of the structure of space-time, to supersymmetry in the fermion case or higher dimensions of space in the vector boson case.

It is remarkable that, in all three classes of models, easily identified radiative corrections give contributions to μ^2 with a negative sign, predicting the instability of the Higgs field to condensation [2]. In all three cases, these contributions come from quantum corrections due to partners of the top quark that are predicted by the new symmetries.

The idea that the condensation of the Higgs field has a definite mechanical explanation from quantum physics thus has major implications. It requires a new set of particles at the TeV mass scale, including exotic partners of the top quark that are expected to be produced at the Large Hadron Collider. This TeV particle spectroscopy can also supply explanations for other issues that require physics beyond the SM. TeV particle spectra typically contain a massive neutral particle that can be absolutely stable and thus a candidate for the particle of dark matter [3]. New couplings among the TeV particles potentially provide new sources of CP violation, offering mechanisms for creating the matter-antimatter asymmetry of the universe. Corrections to the SM coupling constants from the new particles can correct the evolution of the SM couplings, allowing the three SM gauge interactions to unify at very short distances [4].

Most importantly, if the explanation for Higgs condensation changes our view of the SM itself — by making SM particles composite or by enlarging the structure of space-time — these changes must be taken into account in any explanation of phenomena that occur at still smaller distance scales, including the generation of neutrino masses, generation of flavor mixing among quarks and leptons, and the unification of the particle physics interactions with gravity.

In short: mechanisms which shed light on the physics behind the otherwise mysterious potential in Eq. 1.1 are needed to directly address the major experimental anomalies of Section 1.1!

1.2.2 Naturalness

So the major question is: Are there any hints that suggest to us how high in energy must we probe to discover the particles that address the questions of physics beyond the SM? A crisp answer is not yet clear, but we do have a bothersome hint namely from the slippery principle called “naturalness.”

Naturalness is the statement that new particles that generate the μ^2 term in the Higgs potential (1.1) must have masses at the scale of μ^2 itself,

$$\mu^2 \sim (100 \text{ GeV})^2 . \quad (1.3)$$

Taken most naively, naturalness implies that new particles associated with the Higgs potential should have been found in the 1990’s at the experiments at LEP and the Tevatron. Today, the LHC experiments have carried out much deeper searches for these particles. How much further must we go?

One approach to naturalness looks more critically at the radiative corrections to the μ^2 parameter in the SM. The first-order corrections due to the top quark, the W and Z bosons, and the Higgs boson itself are

$$\delta\mu^2 = -\frac{3g_{Htt}^2}{8\pi^2}\Lambda^2 + \frac{3\alpha_w(3 + \tan^2\theta_w)}{4\pi}\Lambda^2 + \frac{\lambda}{8\pi^2}\Lambda^2 , \quad (1.4)$$

where g_{Htt} is the same Yukawa coupling in (1.2), α_w and λ are the couplings of these particles, and θ_w is the weak mixing angle. All three terms are divergent, proportional to Λ^2 , where Λ is a mass scale at which the SM is replaced by a more complete underlying theory. Contributions from new particles add to (or subtract from) this expression. To give a well-defined result for μ^2 , they must cancel the dependence on Λ . If we allow the new contributions to cancel the SM ones over many decimal places, Λ can be arbitrarily high. However, this might be considered “unnatural.” If we assume that at most one significant figure is cancelled, we obtain interesting limits on top, W , and Higgs partners at roughly 2 TeV.

Another approach looks into the computation of μ^2 in specific models [5]. In supersymmetry (SUSY) models, the parameter called μ — the Higgsino mass term — contributes to the Higgs parameter μ^2 at the tree level. Forbidding cancellations beyond one significant figure gives for the SUSY parameter $\mu < 200$ GeV. This is a strong upper bound on the mass of the supersymmetric partner of the Higgs boson, a particle that will be difficult to discover at the LHC. The supersymmetric partners of the top quark and the gluino contribute to μ^2 in one-loop and two-loop order, respectively. The corresponding naturalness bounds are

$$m(\tilde{t}) < 1 \text{ TeV} , \quad m(\tilde{g}) < 2 \text{ TeV} . \quad (1.5)$$

In Little Higgs models in which the Higgs boson is a composite Goldstone boson, the formula for the radiative correction to μ from a new fermionic partner T of the top quark has the form

$$\delta\mu^2 = C \frac{3\lambda_t^2}{8\pi^2} m_T^2 , \quad (1.6)$$

where C is a model-dependent constant of order 1. This gives a bound

$$m(T) < 2 \text{ TeV} . \quad (1.7)$$

In all cases, we might have stronger cancellations in the expressions for μ^2 . Perhaps these cancellations might eventually find some physics explanation. However, each factor of 10 in mass above the bounds quoted requires cancellations of another *two* significant figures. Even such an imprecise criterion as naturalness probably limits top quark partners to lie below about 10 TeV.

183 However unsatisfactory these naturalness estimates might be, our interest in these estimates remains very
184 strong. Higgs condensation is the mechanism that generates the whole spectrum of masses of the SM quarks,
185 leptons, and vector bosons. Can it be just an accident? If not, there *must* be a spectrum of new particles
186 at the TeV scale. Even if we cannot predict the value of this scale incisively, the importance of mass scale
187 is clear. We must find these new states.

188 To investigate new physics at the TeV mass scale, the research program is clear. It consists of three ambitious
189 threads:

- 190 1. First, we must study the properties of the Higgs boson in as much detail as possible.
- 191 2. Second, we must search for the imprint of the TeV mass particles on the heaviest SM particles, the W
192 and Z bosons, and the top quark.
- 193 3. Third, we must search for the direct production of the new particles at high energies.

194 To the extent that the naturalness arguments above are a guide, all three approaches will be accessible at
195 high-energy collider experiments in the near future. In the next section, we will describe the tools that we
196 have available for that search.

197 **1.3 Organization of the Energy Frontier study**

198 In this section, we briefly describe how the Energy Frontier study was organized, in terms of topical working
199 groups and the landscape of proposed accelerators.

200 **1.3.1 Working groups for the study of the Energy Frontier**

201 We divided the study of the TeV energy scale thematically, in terms of probes of this scale using different
202 particles and interactions. The results summarized here constitute the efforts of hundreds of physicists who
203 worked through the winter and spring of 2013 within six working groups. The leaders of these groups are
204 co-authors of this report. The working groups were;

- 205 1. The Higgs Boson
- 206 2. Electroweak Interactions
- 207 3. Quantum Chromodynamics and the Strong Force
- 208 4. Understanding the Top Quark
- 209 5. The Path Beyond the Standard Model - New Particles, Forces, and Dimensions
- 210 6. Flavor Mixing and CP Violation at High Energy

211 Highlights of each group’s work are presented in this order in the following six sections. For each group, the
212 summary of results is followed by their “Message,” a quick summary of their conclusions. We follow with the
213 scientific cases to be made for each possible accelerator organized around each physics group’s conclusions
214 for that facility.

1.3.2 Accelerators for the Study of the Energy Frontier

In our discussion, specific estimates of the capabilities of the methods that we discussed will be made in the context of proposed accelerator programs discussed at Snowmass. We provide here a brief orientation to these programs. Energies refer to the center of mass energy of colliding beam experiments. For details on the design and current status of these proposals, see the Capabilities Frontier working group report [6].

The baseline for our study is Large Hadron Collider (LHC), the pp collider now operating at CERN. The LHC schedule calls for 75-100 fb^{-1} starting in 2015 with, essentially, the current detectors. Following Long Shutdown 2 in approximately 2019, the Phase 1 detector upgrades will be installed and running will resume at a projected instantaneous luminosity of $2 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$. Then, in approximately 2023 the luminosity is expected to increase to $5 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$. In this study we compare the current results from the LHC, at 7-8 TeV with an integrated luminosity of 20 fb^{-1} , to future data samples at 14 TeV with 300 fb^{-1} and with 3000 fb^{-1} . The latter program will often be referred to as the high-luminosity LHC or HL-LHC. The projected evolution of the LHC program is described in [7].

Our study considered higher energy pp colliders, with energy 33 TeV and 100 TeV. Unless it is indicated otherwise, the event sample assumed is 3000 fb^{-1} . A high-energy upgrade of the LHC at 33 TeV (HE-LHC) is discussed in [8]. Colliders of 100 TeV energy are described in [9, 10]. In the following we will refer to such a collider generically as VLHC.

Our study considered e^+e^- linear colliders, both the International Linear Collider (ILC), covering the energy range 90 GeV–1000 GeV and the Compact Linear Collider (CLIC), covering the energy range 350 GeV–3000 GeV. The ILC is described in [11] and in its technical design report [12]. The TDR/CDR luminosity samples are 1000 fb^{-1} at 1 TeV and scaling linearly with energy. Luminosity upgrades of the baseline ILC using strategies outlined in the TDR, to 2500 fb^{-1} at 1 TeV and similar enhancements at other energies, with long running periods, are described in [13]. CLIC is described in [14] and in its Conceptual Design Report [15].

Our study considered $\mu^+\mu^-$ colliders operating over a range from 125 GeV to 3000 GeV. The luminosity samples assumed were similar to those for linear e^+e^- colliders. The technology of the muon collider is described in [16, 17].

Our study considered a circular e^+e^- collider in a large (80-100 km) tunnel. Accelerator parameters for such a machine are described in [18] in the context of one such proposal, TLEP, for a large tunnel near CERN. In principle, accelerator techniques invented for super-B-factors can produce very high luminosities, in excess of $10^{36}/\text{cm}^2\text{sec}$ at 90 GeV and $10^{35}/\text{cm}^2\text{sec}$ at 250 GeV, when summed over 4 detectors. However, there is as yet no complete accelerator design. These machines cannot operate above 350 GeV. In the following, we will refer to such a collider as TLEP. We will assume the above luminosities and operation with four detectors.

Two more types of accelerators received more limited attention from our study. Linear e^+e^- colliders can be converted to photon-photon colliders, with roughly 80% of the energy and similar luminosity, by backscattering laser light from the electron beams. Proposals for photon-photon colliders are described in [19, 20]. Colliding the LHC beam with an e^- or e^+ beam from a linear accelerator offers the opportunity of high energy ep collisions. This has been studied for a facility at CERN called LHeC, described in [21].

1.4 The Higgs Boson

256 We begin with the study of the Higgs boson itself. In this section, we will refer to the new boson with mass
 257 125 GeV as “the Higgs boson,” while recognizing that its properties could well be very different from the
 258 simplest expectations.

259 We have already emphasized that the study of the Higgs boson gives a completely new avenue along which to
 260 probe the physics of the TeV scale. The picture of the Higgs boson given by the SM is precise. All properties
 261 of the Higgs boson can be computed now that the mass of the Higgs boson is known. And yet, this precise
 262 theory has no conceptual foundation. Current experiments exclude deviations from the SM at the 100%
 263 level, but surprises at the 30%, 10%, or 3% level are all possible in different highly plausible models. The
 264 nature of the Higgs boson is a central part of the mystery of TeV physics. New physics responsible for Higgs
 265 condensation must couple to the Higgs boson and affect its properties at some level.

266 Full details of the future program on the Higgs boson, and more precise statements of the uncertainty
 267 estimates given below, can be found in the Higgs Boson working group report [22].

268 1.4.1 Higgs boson couplings

269 The most direct question to ask about the new boson is that of whether it is in fact the sole source of mass
 270 for all quarks, leptons, and gauge bosons. For this to be true, the couplings of the Higgs boson to the various
 271 species of SM particles must follow a definite pattern. The couplings of the particle to fermions and vector
 272 bosons must be **from Eq. 1.2**

$$g_{Hf\bar{f},SM} = \frac{1}{\sqrt{2}} \frac{m_f}{v}, \quad g_{HVV,SM} = \frac{1}{2} \frac{m_V^2}{v^2}, \quad (1.8)$$

273 where v is the value of the SM Higgs condensate, equal to 246 GeV. (More properly; this is the leading-order
 274 prediction for a coupling defined to all orders with all three particles on shell.) These couplings have a simple
 275 pattern that should be tested for as many SM species as possible. In the following discussion, we define

$$\kappa_A = g_{HA\bar{A}}/(SM), \quad (1.9)$$

276 where (SM) denotes the SM prediction.

277 The Higgs boson also couples to pairs of vector bosons gg , $\gamma\gamma$, and γZ through loop diagrams. In the SM,
 278 these couplings are dominated by contributions from W boson and top quark loops. In more general theories,
 279 these couplings can also receive contributions from radiative corrections with new particles in loops. We will
 280 denote ratios of the on-shell couplings to the SM predictions by κ_g , κ_γ , $\kappa_{\gamma Z}$.

281 Corrections to the predictions (1.8) can appear at many levels. If there are multiple Higgs fields that mix
 282 into the observed boson, the κ_A will contain cosines of the mixing angles. These can be as large as the data
 283 **permit. permits.** Radiative corrections to due to loop effects of new particles are expected to be below the
 284 10% level.

285 Corrections to the Higgs couplings are also affected by the Decoupling Theorem [23]: If all new particles have
 286 masses greater than M , we can integrate out these particles. The result is the SM, in which the properties of
 287 the Higgs boson are predicted precisely in terms of its mass. The corrections to the SM values are generated,
 288 in this way of constructing the formalism, by effective higher-dimension operators added to the SM. These
 289 corrections will then be of the order of m_h^2/M^2 . The Decoupling Theorem implies an apparently paradoxical
 290 but nevertheless important conclusion: In a model in which the Higgs sector is very complex but all new

291 particles in it are heavier than 500 GeV, corrections to the Higgs boson properties are at most at the 5-10%
 292 level. We are likely to be in this situation, in which the picture of the Higgs boson is very different from
 293 that in the SM but, since the other particles in the sector are heavy, it is difficult to find this out except by
 294 precision measurement.

295 Typical sizes of Higgs boson coupling modifications are shown in Table 1-1. More details of these estimates
 296 are given in [22].

Model	κ_V	κ_b	κ_γ
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
Decoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$< 1.5\%$
Composite	$\sim -3\%$	$\sim -(3-9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$

Table 1-1. Generic size of Higgs coupling modifications from the Standard Model values when all new particles are $M \sim 1$ TeV and mixing angles satisfy precision electroweak fits.

297 Tests of the values of the Higgs couplings relative to the SM must take account of the theoretical uncertainty
 298 in the comparison to the SM predictions. A potentially observable quantity is the partial decay width
 299 $\Gamma(h \rightarrow A\bar{A})$, related to κ_A by

$$\kappa_A^2 = \Gamma(h \rightarrow A\bar{A}) / (SM) . \quad (1.10)$$

300 Currently, some of these quantities have large uncertainty in their evaluation in the SM. The uncertainty in
 301 the partial width $\Gamma(h \rightarrow b\bar{b})$, which accounts for more than half of the SM Higgs total width, is quoted as 6%
 302 [24]. A concerted program is required to bring the uncertainties in the SM predictions below 1%. This requires
 303 complete evaluation of the 2-loop electroweak corrections to the partial widths. It also requires improvement
 304 of the uncertainty in the crucial input parameters α_s , m_b , and m_c . Lattice gauge theory promises to reduce
 305 the errors on all three quantities to the required levels [25]. Further methods for improvement in our
 306 knowledge of α_s are discussed in Sec. 1.6.

307 There are only a few cases in which the partial widths $\Gamma(h \rightarrow A\bar{A})$ can be measured directly. More often,
 308 the Higgs decay partial widths are measured from the rates of reactions that involve the Higgs boson in an
 309 intermediate state. An example is the rate of $\gamma\gamma$ production through gg fusion at the LHC. The rate of this
 310 process is proportional to

$$\sigma(gg \rightarrow h) \cdot BR(h \rightarrow \gamma\gamma) \sim \frac{\Gamma(h \rightarrow gg)\Gamma(h \rightarrow \gamma\gamma)}{\Gamma_T(h)} , \quad (1.11)$$

311 where $\Gamma_T(h)$ is the total width. In terms of the κ_A quantities, the measured rates are proportional to

$$\sigma(A\bar{A} \rightarrow h)BR(h \rightarrow B\bar{B}) / (SM) = \frac{\kappa_A^2 \kappa_B^2}{\sum_C \kappa_C^2 BR_{SM}(h \rightarrow C\bar{C})} . \quad (1.12)$$

312 The SM prediction for the total width of the Higgs boson is 4 MeV, a value too small to be measured directly
 313 except at a muon collider where the Higgs boson can be produced as a resonance. At all other cases of hadron
 314 and lepton colliders, the total width must be determined by a fit to the collection of measured rates. Such
 315 fits entail some model-dependence to control the size of modes of Higgs decay that are not directly observed.

316 The report [22] compares the capabilities of the LHC and a variety of lepton colliders to extract the values
 317 of the Higgs boson couplings. At the LHC, the total number of Higgs bosons produced is very high, over 170




Figure 1-1. Measurement precision on κ_γ at different facilities.




Figure 1-2. Measurement precision on κ_W at different facilities.




Figure 1-3. Measurement precision on κ_b at different facilities.




Figure 1-4. Measurement precision on κ_t at different facilities.

318 million per experiment for integrated luminosity of 3000 fb^{-1} . However, Higgs boson production at the LHC
 319 is accompanied by very high backgrounds. The extraction of couplings from cross sections is complicated
 320 by significant QCD uncertainties in the calculation of cross sections, currently about 12% for gluon fusion
 321 though smaller for other Higgs production processes.

322 At electron colliders, the Higgs boson is produced in the relatively background-free interactions $\ell^+\ell^- \rightarrow Zh$
 323 and $\ell^+\ell^- \rightarrow \nu\bar{\nu}h$ (vector boson fusion). The measurements of rates are dominated by statistical errors, but
 324 the statistics are limited. The Zh reaction offers tagged Higgs bosons, giving the possibility of observing
 325 invisible and exotic modes of Higgs decay. The total cross section for the two e^+e^- reactions are directly
 326 proportional to $\Gamma(h \rightarrow ZZ^*)$ and $\Gamma(h \rightarrow WW^*)$, respectively, without dependence on the $\Gamma_T(h)$. This allows
 327 lepton collider measurements to determine $\Gamma_T(h)$ by fitting of Higgs rates without any model assumptions.

328 Figures 1-1 through 1-3 show the comparison for a variety of accelerator programs of the projected
 329 uncertainty of measurement of the Higgs couplings to $\gamma\gamma$, WW , and $b\bar{b}$ from an appropriate 6-parameter
 330 fit, and the projected error on the Higgs coupling to $t\bar{t}$ from experiments directly sensitive to this quantity.
 331 The facilities considered are the LHC at its current stage, after 300 fb^{-1} , and after 3000 fb^{-1} , the ILC up
 332 to 500 GeV and up to 1000 GeV, the TLEP circular e^+e^- collider and the CLIC linear collider operating
 333 at 1400 and 3000 GeV. For LHC, the upper and lower estimates reflect a pessimistic scenario, in which
 334 current systematic errors do not improve, and an optimistic scenario in which theory errors are halved and
 335 experimental systematic errors decrease as the square root of the integrated luminosity, For the ILC stages,
 336 the first error bar corresponds to the baseline event samples considered in the ILC TDR, while the second

337 includes, more optimistically, a set of luminosity upgrades described in the TDR. Full details, and tables
 338 of the numerical results of the fit, can be found in [22]. The figures show, first, that the LHC, especially
 339 in its high luminosity phase, will measure Higgs couplings with impressively high precision. However, the
 340 discovery of perturbations of the Higgs boson couplings at the level shown in Table 1-1, at 5σ significance,
 341 will require both the much lower level of systematic errors available at a lepton collider and very large event
 342 samples to reduce the statistical errors.

343 1.4.2 Higgs boson self-coupling

344 A particularly important coupling of the Higgs boson is the Higgs self-coupling, λ in (1.1), which determines
 345 the shape of the Higgs potential. In the SM, after Higgs condensation, there is a triple Higgs boson coupling
 346 proportional to $\sqrt{\lambda}$, given alternatively by

$$\lambda_{hhh} = \frac{3m_h^2}{v^2} . \quad (1.13)$$

347 This coupling can be extracted from the rate for double Higgs production, for example $pp \rightarrow hh + X$ or
 348 $e^+e^- \rightarrow \nu\bar{\nu}hh$.

349 Theoretical models with extended Higgs sectors or Higgs compositeness can predict deviations of the triple
 350 Higgs coupling of 20% relative to the SM expectation. These are larger effects than those expected in the
 351 Higgs couplings to fermions and vector bosons, but the measurement is also much more difficult. The cross
 352 sections at lepton colliders are at the fb level. At the LHC, the cross sections are larger, but the use of
 353 rare decay modes, including $h \rightarrow \gamma\gamma$, is considered to reduce background. Projected uncertainties on λ_{hhh}
 354 are 50% per experiment at the HL-LHC and 13% in a long-term program at the ILC at 1 TeV or 10% for
 355 CLIC at 3 TeV. The double Higgs production cross section increases rapidly with energy. Measurements at
 356 a 100 TeV pp collider are estimated to reach an uncertainty of 8%.

357 1.4.3 Higgs boson spin and CP

358 A crucial test of the identification of the 125 GeV resonance with the Higgs boson is the measurement of
 359 its spin and parity. This issue is almost settled with the current data from the LHC. The fact that the
 360 resonance decays to $\gamma\gamma$ implies that it has integer spin and cannot have spin 1. The distribution of the four
 361 leptons in $h \rightarrow ZZ^*$ decays already strongly favors the 0^+ over the 0^- spin-parity hypothesis and excludes
 362 the simplest forms of spin 2 coupling [26]. This issue should be decided with the next LHC data set.

363 However, there is a more subtle issue associated with the Higgs boson CP. If there are multiple Higgs bosons
 364 and CP violation in the Higgs sector, the Higgs boson at 125 GeV can contain an admixture of CP scalar
 365 states. CP violation in the Higgs sector has major implications. Most importantly, it can provide the new
 366 source of CP violation outside the SM that allows the matter-antimatter asymmetry of the universe to be
 367 generated at the electroweak phase transition.

368 CP violation in the Higgs sector can be reflected both in production and decay of the Higgs boson. The most
 369 accurate tests are available in the study of the 4-lepton final state in $h \rightarrow ZZ^*$. However, CP violating terms
 370 in this vertex can be masked by the large tree level decay amplitude proportional to the Higgs condensate
 371 v . Lepton colliders can search for CP violation in the decay $h \rightarrow \tau^+\tau^-$ and in the production process
 372 $\ell^+\ell^- \rightarrow t\bar{t}h$, reaching 1% precision. Photon-photon colliders, which produce the Higgs boson as a resonance,

373 can use initial-state polarization to search for CP-violating terms in the Higgs boson coupling to $\gamma\gamma$, which
 374 has no tree-level contributions. Similarly, a muon collider can probe for CP-violating contributions to the
 375 Higgs boson coupling to $\mu^+\mu^-$ if the accelerator provides transverse beam polarization.

376 1.4.4 Higgs boson mass and width

377 The Higgs boson mass is currently known from the LHC experiments to better than $600 \text{ MeV}/c^2$. This
 378 accuracy is already sufficient for the uncertainty in the Higgs mass not to be significant in precision
 379 electroweak tests. The most important influence of a highly accurate Higgs mass within the SM comes
 380 in the evaluation of the predictions for the Higgs couplings to WW and ZZ , for which one boson must be
 381 off the mass shell. A $100 \text{ MeV}/c^2$ error in the Higgs mass corresponds to a 0.5% uncertainty in κ_W . We
 382 expect that the error in the Higgs mass can be decreased to $100 \text{ MeV}/c^2$ and to $50 \text{ MeV}/c^2$, respectively,
 383 for the LHC programs with 300 fb^{-1} and 3000 fb^{-1} by using the $\gamma\gamma$, ZZ^* , and $\mu^+\mu^-$ modes in which the
 384 Higgs boson can be fully reconstructed. A lepton collider studying the Higgs boson in the Zh production
 385 mode would push this uncertainty down further, to about $35 \text{ MeV}/c^2$ for linear colliders and $7 \text{ MeV}/c^2$ for
 386 a very high luminosity program at a circular collider.

387 Predictions of the Higgs mass in models of new physics might provide further motivation for measuring
 388 the Higgs mass accurately. An example of such a model is the Minimal Supersymmetric Standard Model
 389 (MSSM). To evaluate the prediction to an accuracy of $100 \text{ MeV}/c^2$, however, the masses of the top squarks
 390 must be known, and the top quark mass must be known to $100 \text{ MeV}/c^2$.

391 We have noted already that lepton colliders offer the possibility of a model-independent determination of
 392 the Higgs boson total width. Because the couplings of the Higgs boson to ZZ and WW appear both in the
 393 expressions for measurable total cross sections and branching ratios, these couplings can be eliminated to
 394 evaluate the total width through the relations

$$\Gamma_T(h) \sim \sigma(\ell^+\ell^- \rightarrow Zh)/BR(h \rightarrow ZZ^*) \sim \sigma(\ell^+\ell^- \rightarrow \nu\bar{\nu}h, h \rightarrow b\bar{b})/BR(h \rightarrow WW^*)BR(h \rightarrow b\bar{b}) \quad (1.14)$$

395 This gives the Higgs boson width to 3% for a long-term program at the ILC and to 0.6% for a high luminosity
 396 program at a circular collider with multiple detectors. These uncertainties are reflected in the coupling
 397 uncertainties quoted in Sec. 1.4.1.

398 A muon collider would have the capability of observing the Higgs boson as a narrow resonance. For the
 399 projected beam energy resolution of 4×10^{-5} , the mass of the Higgs boson would be measured to 0.06 MeV
 400 and the width would be measured directly **in the s -channel** to a precision of 4% [17]

401 1.4.5 Searches for additional Higgs bosons

402 There are strong motivations for expecting the existence of additional Higgs particles. These motivations
 403 begin with the overall mysteries of the physics of Higgs condensation and the question of whether the Higgs
 404 **boson** is the only particle of the SM whose quantum numbers do not come in multiples. Beyond this,
 405 virtually all models of new physics to explain the Higgs potential contained additional Higgs doublet fields.
 406 These fields are required in supersymmetric models in order for Higgs fields give mass to both the up-type
 407 and the down-type quarks. In models with new space dimensions, additional Higgs fields arise as the Kaluza-
 408 Klein excitations of the fundamental Higgs doublet. Each additional Higgs doublet gives rise to four new
 409 particles, CP-even and CP-odd neutral scalars H and A , and a charged pair H^\pm .

410 A typical feature of additional Higgs particles is that the associated fields have much smaller couplings to
 411 WW and ZZ than for the lightest Higgs boson. Often, these particles have enhanced couplings to heavy
 412 flavors, either to b and τ or to t depending on whether the extended Higgs parameter $\tan\beta$ is greater than
 413 or less than 1. This de-emphasizes searches based on vector boson fusion in favor of search techniques that
 414 involve $b\bar{b}$ annihilation to Higgs resonances.

415 Currently, the LHC experiments exclude additional Higgs bosons for masses as high as 1 TeV in restricted
 416 ranges of $\tan\beta$. The region of large $\tan\beta$ is surveyed by reactions such as $b\bar{b} \rightarrow H, A \rightarrow \tau^+\tau^-$, while the
 417 region of low $\tan\beta$ is surveyed by reactions such as $gg \rightarrow H, A \rightarrow t\bar{t}, gg \rightarrow A \rightarrow Zh$. A gap remains for
 418 intermediate values, roughly $2 < \tan\beta < 20$, which is closed only for extended Higgs boson masses up to
 419 200 GeV. Future runs of the LHC, up to 3000 fb^{-1} , are expected to close this window up to about 500 GeV.

420 Lepton collider experiments can search for extended Higgs boson states through the reaction $\ell^+\ell^- \rightarrow HA$ up
 421 to the kinematic limit, independently of the value of $\tan\beta$. This covers the parameter space up to 500 GeV
 422 for ILC and up to 1500 GeV for CLIC running at 3 TeV. Photon and muon colliders have the opportunity
 423 to discover additional Higgs bosons as resonances up to the full center of mass energy of the machine.

424 1.4.6 The Message

425 The conclusions of the Higgs Boson working group can be summarized as follows:

- 426 1. Direct measurement of the Higgs boson is the key to understanding electroweak symmetry breaking.
 427 The fact that the Higgs boson appears as a light, apparently fundamental, scalar particle needs
 428 explanation. A research program focused on the Higgs couplings to fermions and vector bosons and
 429 achieving a precision of a few percent or less is required to address these questions.
- 430 2. Full exploitation of the LHC is the path to few percent precision in the Higgs coupling and to a 50
 431 MeV precision in the determination of the Higgs mass.
- 432 3. Full exploitation of a precision electron collider is the path to a model-independent measurement of
 433 the Higgs boson width and a sub-percent measurement of the Higgs couplings, allowing discovery of
 434 new physics effects at the percent level.

435 Experiments on Higgs bosons give information on the Particle Physics Questions # 1, 2, 4, 5, 8, 9, 10 listed
 436 in the Snowmass Summary [27].

438 1.5 Electroweak Interactions

440 The precision electroweak experiments of the 1990's established the $SU(2) \times U(1)$ theory of electroweak
 441 interactions at the sub-percent level of accuracy. But, they did more. All particle species with couplings
 442 to the electroweak interactions eventually influence the properties of the weak interaction bosons, W and
 443 Z . Very precise measurements of the properties of these boson then have the potential to reveal new,
 444 undiscovered particles. The experiments of the 1990's indicated the presence of a heavy top quark and
 445 a light Higgs boson and estimated the masses at which these particles were eventually discovered. They
 446 disfavored a fourth generation of quarks and leptons, now excluded by direct search at the LHC.

447 Increased precision in the properties of the weak interaction bosons could well turn up the first evidence of
 448 the TeV spectrum of particles discussed in Sec. 1.2.1. There still tensions evident in the data which have
 449 been intriguing for years; for example, the current value of M_W from many experiments persists at about
 450 1σ higher than the SM expectation. Experiments over the next decade will explore whether these could
 451 become significant deviations requiring radiative corrections due to new particles.

452 In a description of possible new interactions in terms of effective operators, the electroweak precision
 453 observables probe only the first few terms. Experiments at higher energy probe additional operators by
 454 observing and constraining the nonlinear interactions of the W and Z . These operators can receive corrections
 455 from loop diagrams involving new TeV mass particles but, more strikingly, they can receive leading-order
 456 corrections if there is new strong dynamics or resonances in the Higgs sector.

457 We will review these topics in this section. Full details of the program, and more precise statements of the
 458 projected uncertainties described below, can be found in the Electroweak Interactions working group report
 459 [28].

460 1.5.1 Precision observables M_W and $\sin^2\theta_w$

461 The current uncertainty in the W boson mass is 15 MeV, corresponding to an relative precision of 2×10^{-4} .
 462 It is remarkable that the most accurate determinations of M_W come from the hadron collider experiments
 463 CDF and DO.

464 Precision measurement of M_W at hadron colliders is very challenging, but certain features of W production
 465 make it feasible to reach high accuracies. The directly measured transverse mass distribution is very sensitive
 466 to M_W , having a relatively sharp endpoint at the W mass. Likewise the p_T distributions of the leptons are
 467 also sensitive to the boson mass with different, but manageable systematic uncertainties. Enormous statistics
 468 will be available with very small contamination by background. The dominant errors come from the small
 469 corrections to these properties. Currently, the largest source of systematic error is the dependence of the
 470 acceptance on the rapidity of the produced W , requiring a correction that depends on quark and antiquark
 471 parton distribution functions (PDFs). Experimental uncertainties are at the same level as those due to PDFs
 472 and are expected to continue to decrease accordingly.

473 We see good prospects for improving this measurement at the LHC. The statistical component of the error
 474 will be negligible already with the current LHC data set. The error from PDFs doubles in going from the
 475 Tevatron to the LHC because proton-proton collisions give no valence antiquarks. However, we anticipate
 476 that this error will be decreased using new data on the vector boson rapidity and charge asymmetries. The
 477 issue of PDF improvement is discussed further in Sec. 1.6.2. The huge statistical precision will allow for
 478 control of calorimetric and tracking systematic uncertainties. In Table 1-2 we see that the PDF error in
 479 M_W can be brought down to ± 5 MeV with 300 fb^{-1} and to ± 3 MeV with 3000 fb^{-1} , leading to a final
 480 uncertainty in M_W of ± 5 MeV. In each stage, the experimental systematics are expected to keep pace. In
 481 order to reach this important level of precision, the PDF uncertainties must be pushed to a factor of 7 better
 482 precision that currently available.

483 Lepton colliders offer an opportunity to push the uncertainty in M_W down even further. The W mass was
 484 measured at LEP to ± 36 MeV from the kinematics of W^+W^- production. The uncertainty was dominated
 485 by statistical errors, with a substantial addition contribution from the modeling of hadronization. Both
 486 sources will benefit from the data set on W^+W^- , about 1000 times larger, that will be available at next-
 487 generation e^+e^- colliders such as ILC and TLEP. We estimate an error below ± 4 MeV from this method,
 488 and a similar error from independent measurements on single W production.

ΔM_W [MeV]	LHC		
\sqrt{s} [TeV]	8	14	14
\mathcal{L} [fb ⁻¹]	20	300	3000
PDF	10	5	3
QED rad.	4	3	2
$p_T(W)$ model	2	1	1
other systematics	10	5	3
W statistics	1	0.2	0
Total	15	8	5

Table 1-2. Current and target uncertainties in the measurement of M_W at the LHC.

489 The ultimate W mass measurement would come from a dedicated energy scan of the W^+W^- threshold at
 490 160 GeV. Such a measurement could reach ± 2.5 MeV with the statistics available from the ILC and \pm
 491 1 MeV with the statistics available from TLEP. At this level, systematic errors become dominant. The
 492 program also requires a detailed precision theory of the W^+W^- threshold, using methods now applied to
 493 the $t\bar{t}$ threshold.

494 The measurement of the value of $\sin^2\theta_w$ associated with quark and lepton couplings to the Z resonance
 495 offers an orthogonal probe of the electroweak interactions. The current accuracy in $\sin^2\theta_w$ is at the $7 \times$
 496 10^{-5} level of precision and ~~is~~ is dominated by measurements from LEP and SLC. This level might be
 497 reached but, we expect, will not be surpassed at the LHC. Again, uncertainties in PDFs give the limiting
 498 systematic error. Measurements from the polarization-dependence of the Z cross section and from the b
 499 quark forward-backward asymmetry are discrepant by about 3σ , indicating an experimental question that
 500 should be resolved.

501 Future lepton colliders give an opportunity to improve the value of $\sin^2\theta_w$. The ILC program includes a few
 502 months of running at the Z resonance to produce a data set of 10^9 Z 's, improving the statistics from LEP
 503 by a factor of 100 with highly polarized beams. The ILC detectors should also dramatically improve the
 504 capability for heavy flavor tagging. This "Giga- Z " program should improve the uncertainty in $\sin^2\theta_w$ by a
 505 factor 10. The program also would give new measurements of other Z pole observables sensitive to new TeV
 506 mass particles, most importantly, the fraction R_b of Z decays to $b\bar{b}$.

507 TLEP envisions a multi-year program at higher luminosity to collect 10^{12} events on the Z resonance.
 508 This potentially pushes the precision of electroweak measurements by another order of magnitude, though
 509 systematic contributions to the errors must still be understood. Among other factors, the Z mass must be
 510 measured more accurately than the current 2.5 MeV. This is possible at TLEP if transverse polarization can
 511 be achieved in single beam operation. The direct measurement of $\sin^2\theta_w$ optimally requires longitudinal
 512 polarization in colliding beam mode; the feasibility of this at TLEP needs to be understood.

513 Loop effects from TeV mass particles can produce effects at the 10^{-4} level in both M_W and $\sin^2\theta_w$, so
 514 the improved capabilities for precision electroweak may point to new particle discovery or confirmation.
 515 Quantitative estimates for a number of models are given in [28].

516 1.5.2 Interactions of W and Z bosons

517 The interactions of W and Z bosons are studied at higher energies, through the measurement of vector
518 boson pair production and multi-vector boson production. This study has already begun at LEP and the
519 Tevatron, where parameters of the 3-gauge boson interactions were bounded within a few percent of their
520 SM values.

521 Vector boson interactions are described in a unified way through the formalism of effective Lagrangians.
522 One begins from the SM Lagrangian, in which the Yang-Mills vertices for γ , W , and Z appear as terms of
523 dimension 4. One then adds higher dimension operators invariant under the $SU(2) \times U(1)$ gauge symmetry.
524 A typical term involving an operator of dimension 6 is

$$\delta\mathcal{L} = \frac{c_W}{\Lambda^2} (D_\mu\Phi)^\dagger W_{\mu\nu} (D_\nu\Phi) , \quad (1.15)$$

525 where Φ is the Higgs doublet field and $W_{\mu\nu}$ is the W boson field strength. This operators contributes to
526 the 3- and 4-vector boson vertices. Additional operators of dimension 8 can modify the 4-vector interactions
527 independently of the 3-vector interactions. A typical term is

$$\delta\mathcal{L} = \frac{f_{T,0}}{\Lambda^4} \text{tr}(W_{\mu\nu})^2 \text{tr}(W_{\lambda\sigma})^2 . \quad (1.16)$$

528 In a weak-coupling theory such as the SM, the coefficients c_i and f_j are induced by loop diagrams and should
529 be highly suppressed, by powers of $\alpha_w/4\pi \sim 10^{-3}$. However, in theories with strong interactions in the Higgs
530 sector, the c_i and f_j coefficients could be of order 1, with the Λ parameters then interpreted as the masses
531 of Higgs sector resonances. For example, the operator (1.16) would be induced by a scalar resonance in the
532 Higgs sector.

533 The current bounds on triple gauge boson couplings imply that the Λ parameters associated with dimension
534 6 operators are higher than about 600 GeV. High statistics measurements of the the triple gauge bosons by
535 observation of W^+W^- and ZZ production in e^+e^- reactions at 500 GeV are expected to be sensitive to
536 deviations from the SM that are 10 times smaller, pushing the sensitivity to Λ almost to 2 TeV.

537 It is difficult for hadron colliders to have similar sensitivity to triple gauge couplings. One source of this
538 difficulty is that that the LHC experiments study diboson reactions at higher energies, where additional
539 terms from higher dimension operators are important and so the extraction of the coefficients of dimension
540 6 operators is model-dependent. Of course, there is a compensatory advantage. Working at higher energy,
541 the LHC will study W and Z bosons at energies where Higgs sector strong interactions can dramatically
542 alter the amplitudes for vector boson scattering.

543 Some quantitative examples are presented in [28]. In examples studied there, the LHC at 300 fb^{-1} would
544 achieve exclusion of a Λ value in dimension 8 operators greater than 1.2 TeV. The HL-LHC, with 3000 fb^{-1} ,
545 would roughly triple the significance of the effect, allowing 5σ discovery at 1.2 TeV or exclusion up to
546 1.5 TeV.

547 Increasing the energy allowed for the diboson system dramatically increases the physics reach. A 33 TeV
548 pp collider would push the discovery reach close to 3 TeV, well into the region in which new Higgs sector
549 dynamics would be expected in models of this type.

550 1.5.3 The Message

551 The conclusions of the Electroweak Interactions working group can be summarized as follows:

- 552 1. Precision measurements of the W and Z bosons has the potential to probe indirectly for new particles
553 with TeV masses. This precision program is within the capabilities of LHC, linear e^+e^- colliders, and
554 TLEP.
- 555 2. Measurement of vector boson interaction will probe for new dynamics in the Higgs sector. In such
556 theories, we expect correlated signals in triple and quartic gauge couplings. The LHC and linear
557 colliders will have sensitivity into the mass region above 1 TeV.

558 Experiments on electroweak interactions give information on the Particle Physics Questions # 1, 4, 8, 9
559 listed in the Snowmass Summary [27].

561 1.6 Quantum Chromodynamics and the Strong Interaction

563 Every probe of particle physics at high energies eventually requires detailed knowledge of the strong inter-
564 actions. Even in pure electroweak processes, the strong interactions affect the values of α and α_w through
565 coupling constant renormalization. In the next decade, when most of our new knowledge about particle
566 physics will come from hadron collider experiments, our understanding of the strong interactions will affect
567 every aspect of the data, through the structure of the proton, through radiative corrections to initial- and
568 final-state quarks and gluons and their transition to hadrons, and through the detailed physics that produces
569 multi-jet events.

570 The strong interactions are known to be described by the Yang-Mills theory Quantum Chromodynamics
571 (QCD). This is a theory that is weakly coupled at short distances and strongly coupled at large distances.
572 Our understanding of QCD is imperfect. We have limited tools for the strongly coupled regime, and precision
573 calculation in the weakly coupled regime is technically complex. Nevertheless, our knowledge of QCD has
574 taken enormous strides since the previous Snowmass workshop a decade ago. In this section, we review the
575 current state of our tools for QCD and indicate the opportunities for further progress. More details on all
576 of the topics discussed here can be found in the working group report [29].

577 In our discussion of precision quantum field theory calculation, we will describe one-loop radiative corrections
578 as next-to-leading order (NLO), and higher corrections as NNLO, *etc.*

579 1.6.1 α_s

580 The strength of the QCD coupling is determined by the value of the coupling constant α_s , usually expressed
581 as its \overline{MS} value at the Z mass. The current Particle Data Group [30] value of this quantity has 0.6%
582 uncertainty. We have pointed out in Sec. 1.4.1 that this is a limiting systematic error in the evaluation of
583 the SM prediction for Higgs boson couplings.

584 There are three strategies to improve the estimate. The extraction of α_s from the Z boson width or the
585 ratio of hadronic to leptonic Z decays is theoretically unambiguous but is limited by statistics. The Giga- Z
586 program at the ILC discussed in the previous section should improve the current determination from this
587 source from 4% to 0.4%. The very high luminosity Z program envisioned for TLEP could decrease this
588 uncertainty further to 0.1%. We judge that higher-statistics measurements of e^+e^- event shapes are not
589 competitive with these improvements.

590 Proposed improvements of PDFs from LHeC will lead to an improved value of α_s from the measurement
 591 of PDF evolution. The expected statistical error would be $\pm 0.2\%$ from LHeC alone and $\pm 0.1\%$ from the
 592 combination of LHeC and HERA. The theoretical systematic error for this method is not as well understood,
 593 but we estimate this at $\pm 0.5\%$ once one further order in QCD perturbation theory (N³LO) is calculated.

594 The most accurate current values of α_s come from lattice gauge theory. Higher statistics lattice estimates
 595 and calculation of additional terms in lattice perturbation theory should decrease the current uncertainties
 596 over the next decade to 0.3%. These improvements will come together with improvements in the values of
 597 the quark masses, as discussed in 1.4.1.

598 1.6.2 Parton distribution functions

599 Our knowledge of the initial state in hadron-hadron collisions is encoded in the representation of the proton
 600 structure given by the parton distribution functions (PDFs). The provision of PDF distributions with
 601 uncertainties is an innovation of the past decade [31]. The uncertainties quoted have been continually
 602 improved through the addition of new data sets, especially from the Tevatron and HERA experiments.




Figure 1-5. TBA need higher resolution version.

603 Still, there are gaps in our knowledge, especially in the components relevant to the study of physics beyond
 604 the SM. The leading PDF distributions disagree in their estimates of the gluon-gluon luminosity function at
 605 the mass of the Higgs boson; this accounts for an 8% systematic uncertainty in the extraction of the cross
 606 section for Higgs production. The step from the Tevatron to the LHC puts added weight on the antiquark
 607 distributions in the proton. And, all parton luminosities are poorly constrained by data for parton-parton
 608 invariant masses greater than about 500 GeV. This is clearly shown in Fig. 1-5.

609 We expect that these difficulties can be address using future data from the LHC. PDFs at the few-percent
 610 level of accuracy require theoretical calculations at the NNLO level. These are already available for the
 611 Drell-Yan process [32]. The NNLO computation of the total cross section for top quark pair production has
 612 recently been completed [33] and is now being extended to the rapidity distribution. NNLO calculations
 613 for 2-jet production are in progress. Over the next few years, these calculations will be used in conjunction
 614 with a very high-statistics data set on jet, top quark, and lepton pair production from the LHC. The LHCb
 615 experiment has an important role to play in the measurement of rapidity distributions at $y > 2.5$ [34].

616 Further improvements in PDFs can result from the program of the LHeC. The data expected will reduce
617 the error in the gluon luminosity to a few percent at the Higgs boson mass and to 5-10% in the multi-TeV
618 region.

619 1.6.3 Electroweak corrections to hadron collider processes

620 The quest for few-percent accuracy in predictions for hadron colliders brings new elements into play. In
621 particular, it requires that QED and electroweak corrections be included in all predictions for LHC.

622 Three elements are needed here. Electroweak corrections at NLO order are generally comparable to NNLO
623 QCD corrections, so both types of corrections should be included, together with $\mathcal{O}(\alpha_w \alpha_s)$ terms if possible.
624 Electromagnetic corrections to hadronic reactions cannot be consistently included without a set of PDFs
625 derived from formulae that include NLO QED corrections. This requires a nontrivial modification of PDF
626 fitting programs in order to introduce a photon PDF for the proton. [Guess I have trouble referring to a photon
627 as a parton.](#) Photon-induced reactions can contribute to LHC processes at the few-percent level, increasing
628 to the 10% level at higher pp energies.

629 **Finally** ~~Finally~~, at energies of a TeV and above, electroweak Sudakov effects, negative corrections to
630 two-particle production proportional to $\alpha_w \log^2 s/M_W^2$, can become important. These are 10% corrections
631 for Drell-Yan processes producing 3 TeV dilepton systems. At higher energy pp colliders, these double
632 logarithmic corrections must be resummed systematically.

633 1.6.4 High-precision calculation

634 In the past decade, a revolution in calculational technique has made it possible to derive formulae at NLO
635 for the QCD cross sections for complex multiparton processes such as $pp \rightarrow W + 4$ jets and $pp \rightarrow t\bar{t} + 2$ jets.
636 This has reduced the size of the theoretical errors in these cross sections from order 1 to 10-20%. Methods
637 are now being developed to evaluate general 2-parton processes and even some 3-parton production processes
638 to NNLO, to reduce these theoretical errors to the few-percent level.

639 We have already made reference to NNLO calculations of $t\bar{t}$ and 2-jet production. A very important target
640 here is the cross section for Higgs boson production in association with one or more jets. Many Higgs
641 measurements at the LHC include jet vetoes to control background from $t\bar{t}$ production and other sources,
642 so explicit accounting for emitted jets is necessary. These cross sections often require terms to NNLO for
643 stable summation of the perturbation series.

644 Beyond the fixed-order perturbation theory, many other aspects of higher-order computation remain to be
645 understood. NNLO computations often display large logarithms, which should be systematically resummed.
646 The merging of Monte Carlo programs with NLO QCD calculations is incompletely understood, and new
647 difficulties arise at NNLO. We are optimistic that Higgs boson production and other QCD processes can be
648 computed to few-percent accuracy, but many challenges remain.

649 1.6.5 The Message

650 The conclusions of the QCD working group can be summarized as follows:

- 651 1. Improvements in PDF uncertainties are required. There are strategies at LHC for these improvements.
652 QED and electroweak corrections must be included in PDFs and in perturbative calculations.
- 653 2. An uncertainty in α_s of order 0.1% may be achievable through improvements in lattice gauge theory
654 and precision experiments.
- 655 3. Advances in all collider experiments, especially on the Higgs boson, require continued advances in
656 perturbative QCD.

657 Experiments on QCD give information on the Particle Physics Questions # 1, 2, 8, 9 listed in the Snowmass
658 Summary [27].

659

660 1.7 Fully Understanding the Top Quark

661

662 The top quark is the heaviest quark and, indeed, the heaviest elementary particle known today. Its large
663 mass give it the strongest coupling to the Higgs boson and to other possible particles of the Higgs sector.
664 The mass of the top quark seems to be anomalously large—though it is sometime argued that it is the masses
665 of all other quarks and leptons that are anomalously small. For all of these reasons, the top quark merits
666 thorough experimental investigation.

667 The Tevatron experiments that discovered the top quark produced, in all, about 100,000 of these particles.
668 The LHC experiments have already produced 10 million and aim for many billions of top quarks by the end
669 of the HL-LHC. Future lepton colliders will bring new precision tools to the study of the top quark. In this
670 section, we will discuss what can be learned from these observations. More details on all of these topics can
671 be found in the working group report [35].

672 1.7.1 Top quark mass

673 Like α_s discussed in the previous section, the top quark mass is a crucial input parameter for many SM
674 predictions. It is already the most accurately known quark mass; a 2 GeV uncertainty on this quantity
675 corresponds to a measurement with 1% precision. An accurate top quark mass is needed for precision
676 electroweak fits, with an error of 600 MeV on the top quark mass yielding, for example, an error of 5 MeV
677 in M_W . The top quark mass is also an important input to the question of ultimate vacuum stability in the
678 SM [36]. Figure 1-6 shows the suspiciously marginal position of the measured Higgs boson mass and that
679 of the top quark. Clearly more precision on the latter would help to elucidate whether this is another hint
680 of strange behavior in the Higgs sector.

681 The top quark mass is most precisely defined as an \overline{MS} quantity, evaluated most conveniently at the \overline{MS} top
682 quark mass value itself. However, experimental determinations of the top quark mass are typically done by
683 kinematic fitting to templates, with poorly controlled errors from Monte Carlo modeling and hadronization.
684 Thus, the precision determination of the top quark mass contains a double challenge, first, to give a definition
685 of the top quark mass that can be cleanly related to the \overline{MS} mass, and then to measure that quantity
686 accurately.




Figure 1-6. Regions of metastability and instability of the Higgs potential in the SM, as the top quark and Higgs boson masses are varied, from [36].

687 One solution to this challenge at the LHC is an idea from CMS[37] to measure the top quark mass from the
 688 endpoint of the distribution of the mass $m(b\ell)$ in top quark pair production, where ℓ is an isolated lepton and
 689 b is a b -tagged jet defined by anti- k_T or a similar prescription. The distribution of $m(b\ell)$ can be computed in
 690 QCD perturbation theory in terms of the perturbative pole mass, which can be related to the \overline{MS} mass with
 691 an error of the order of 200 MeV. The endpoint feature in the distribution is sharp and strongly dependent
 692 on m_t . We expect that this method can reach a total uncertainty of 500 MeV with the statistics of the
 693 HL-LHC. Other methods for measuring the top quark mass at the LHC are discussed in [35].

694 At lepton colliders, the cross section for $t\bar{t}$ production near the threshold has a distinctive rise sensitive to the
 695 position of the lowest (unstable) $t\bar{t}$ bound state. Extensive theoretical work has evaluated this cross section
 696 to NNLO, with resummation of all large logarithms. The threshold position can be measured to 35 MeV at
 697 the ILC and somewhat better at TLEP and muon colliders. The conversion to the \overline{MS} mass gives a total
 698 uncertainty of about 100 MeV. This very accurate value of m_t is well matched to the precision electroweak
 699 programs at lepton colliders described in Sec. 1.5.

700 1.7.2 Strong and electroweak couplings

701 The production and dynamics of top quark pairs at colliders offers many opportunities to test the strong and
 702 electroweak couplings of these particles. At hadron colliders, the dominant pair production mechanism is
 703 through QCD. The current agreement between the predicted and measured values verifies that the absolute
 704 strength of the QCD coupling to the top quark is equal to the value of α_s measured elsewhere to about 3%
 705 accuracy.

706 Changes in the form of the top quark coupling to gluons might be induced by new resonances associated with
 707 top quark compositeness. Possible magnetic or electric dipole couplings can be probed from the kinematics
 708 of top final states to better than 1% at the LHC with 300 fb^{-1} . Though it is difficult to measure the absolute
 709 size of the top quark width at a hadron collider, the W helicity fractions in top quark decay are sensitive
 710 to modifications of the top quark coupling to W , with similar sensitivity. The cross section for single top
 711 production provides a measure of the CKM matrix element V_{tb} which should reach an accuracy of 2.5% at
 712 300 fb^{-1} . Couplings of the top quark to the photon and Z are constrained by measurements of radiation
 713 from a $t\bar{t}$ state. The HL-LHC is expected to reach sensitivities of a few percent for the photon couplings and
 714 15-20% for the Z couplings.

At lepton colliders, $t\bar{t}$ pairs are produced through virtual photons and Z s, with large interference effects that depend on the beam polarization. The ILC and CLIC, which can take advantage of large beam polarization, expect to reach sensitivities below the 1% level for both photon and Z couplings. Randall-Sundrum models and other models with Higgs compositeness predict shifts of the Z couplings to $t\bar{t}$ at the few percent level; these effects could potentially be discovered in the linear collider programs.

1.7.3 Rare decays

The large samples of top quarks available at the LHC allow deep searches for flavor-changing top quark decays. Neutral current decays of the top quark such as $t \rightarrow \gamma c$ or $t \rightarrow gc$ are utterly negligible in the SM, with predict branching ratios smaller than 10^{-12} . These decays do appear in models with an extended Higgs sector or R-parity violation supersymmetric couplings that bring in two structures of flavor mixing. Searches for these decays at the HL-LHC can reach branching ratio limits below 10^{-5} .

Lepton colliders can also access these flavor changing couplings in single top production, for example, through $\gamma^*, Z^* \rightarrow t\bar{c}, t\bar{u}$. Searches for these processes can reach sensitivities close to 10^{-4} even in experiments at 250 GeV, below the $t\bar{t}$ threshold, and below 10^{-5} in the full ILC program at 500 GeV.

More details on the specific estimates for each possible neutral current coupling can be found in [35].

1.7.4 Searches for new particles related to the top quark

The motivation that we have given for new particles at the TeV scale in Sec. 1.2.2 directly implies the presence of exotic partners of the top quark. Examples of these particles are scalar top quarks in models of SUSY and Kaluza Klein excitations of top quarks in models with extra space dimensions. Searches for these particles have been a very high priority in the LHC program and will continue to be pursued intensively as more data accumulates.

Searches are designed individually for each type of exotic particle. The most powerful searches make use of the fact that top quarks resulting from the decay of the particle have different polarizations than those typically produced in SM pair-production. This is reflected in the kinematic distributions of the $t\bar{t}$ final states. For particles with masses of 1 TeV and above, the preferred method for identification of final-state top quarks is as exotic jets with a 3-jet substructure [38]. This “boosted top” identification is quite insensitive to the pileup associated with high luminosity.

Top squarks in SUSY might be tied to the general mass scale of supersymmetric particles, but the naturalness arguments we have given in Sec. 1.2.2 indicate that they might be the lightest colored supersymmetric partners. The LHC experiments have searched extensively for direct pair-production of top squarks that decay to the lightest supersymmetric particle $\tilde{\chi}^0$ through $\tilde{t} \rightarrow t\tilde{\chi}^0$ and $\tilde{t} \rightarrow b\tilde{\chi}^+$. Current searches exclude a top squark up to about 650 GeV in the limit of light electroweak superpartners. The sensitivity should advance to about 1 TeV at 14 TeV and 300 fb $^{-1}$, and to 1.2 TeV with 3000 fb $^{-1}$.

In composite Higgs and extra dimensional models, the expected partners of the top quark are fermions with vectorlike couplings. The searches for these particles are similar to those for fourth-generation quarks, but they involve more complex decay patterns, with $T \rightarrow Wb$, $T \rightarrow tZ$ and $T \rightarrow th$. Searches for these particles that are comprehensive with respect to the decay mode currently exclude vectorlike top partners

752 up to masses of about 650 GeV. The 14 TeV stages of the LHC will be able to discover these particles at
753 masses of sensitivity to 1.3 TeV for 300 fb⁻¹ and to 1.6 TeV for 3000 fb⁻¹.

754 Composite Higgs models also typically include resonances in the multi-TeV mass region that decay prefer-
755 entially to $t\bar{t}$. Randall-Sundrum models, for example, predict a resonance at a mass of a few TeV decaying
756 with high top quark polarization to $t_R\bar{t}_L$. The boosted top identification described above was developed
757 for the problem of discovering such states and is indeed expected to be very effective. Applying the same
758 methods to larger data sets, we expect a sensitivity to such resonances up to 4.5 TeV for the 14 TeV LHC
759 with 300 fb⁻¹ and up to 6.5 TeV with 3000 fb⁻¹.

760 Additional examples of new particle searches involving top quarks are described in [35].

761 1.7.5 The Message

762 The conclusions of the Top Quark working group can be summarized as follows:

- 763 1. The top quark is intimately tied to the problems of symmetry breaking and flavor.
- 764 2. Precise and theoretically well-understood measurements of top quark masses are possible both at LHC
765 and at e^+e^- colliders, in each case, matching the needs of the precision electroweak program.
- 766 3. New top couplings and new particles decaying to top play a key role in models of Higgs symmetry
767 breaking. LHC will search for the new particles directly. Linear collider experiments will be sensitive
768 to predicted deviations from the SM in the top quark couplings.

769 Experiments on the top quark give information on the Particle Physics Questions # 1, 2, 4, 8, 9 listed in
770 the Snowmass Summary [27].

771

772 1.8 The Path Beyond the Standard Model - New Particles, Forces, 773 and Dimensions

774

775 Models of new physics associated with the TeV mass scale contain a wide variety of new particles. These
776 include the partners of the top quark discussed in the previous section. More generally, many of the schemes
777 discussed in Sec. 1.2.2 for explaining Higgs condensation are based on far-reaching principles that require
778 a spectroscopy of new particles containing heavy partners for all SM particles. This includes additional
779 strongly interacting particles, particles with only electroweak interactions, and new vector bosons. Some of
780 these particles may have lifetimes long enough that their decays are not prompt in a collider experiment.
781 One or more of these particles could be the constituents of the cosmic dark matter.

782 New particles could also introduce new flavor-changing interactions and this intersect with searches for flavor
783 and CP violation in rare processes. This subject will be discussed in Sec. 1.9.

784 The LHC experiments have been able to search for new particles very robustly over a broad range of
785 properties. In this section, we will discuss how higher energies and luminosities at hadron colliders and new
786 capabilities of lepton colliders will extend these searches.

787 Models of Higgs condensation and other TeV-scale phenomena based on different underlying principles make
788 qualitatively different predictions for the quantum numbers and mass relations of the new particle spectrum.
789 Thus, the first discovery of a new particle beyond the Standard Model will define a direction for an extensive
790 research program, one that will be carried out over decades with multiple, complementary, experiments.
791 In this section, we will emphasize the comparative reach of proposed collider programs to make this first
792 discovery. Examples of the consequences of such a discovery will be given in Sec. 1.11. We will have room
793 to discuss only a limited number of examples. The full range of searches for new particles accessible to TeV
794 energy experiments is described in the New Particles and Forces working group report [39].

795 1.8.1 New Vector Bosons

796 We begin by discussing particles that show up in collider experiments as distinct resonances. An example is a
797 color-singlet vector boson associated with an extension of the Yang-Mills symmetry group beyond that of the
798 SM. Such bosons are required in many contexts, including models with left-right symmetric weak interactions
799 at high energy, models of the Higgsino mass in SUSY, and models with extra dimensions. Models of Higgs
800 composite structure often require breaking of a larger gauge group to the SM symmetry group.

801 Searches for these bosons are conducted at hadron colliders by looking for narrow dilepton resonances. A
802 typical benchmark is sensitivity to the “sequential SM” Z' , a boson with the couplings of the Z but with a
803 higher mass. Current results from the LHC require the mass of such a particle to be above 2.5 TeV. With 14
804 TeV, it will be possible to discover such a resonance at 4.5 TeV, for 300 fb^{-1} , and at 7 TeV, for 3000 fb^{-1} .
805 The value of the production cross section and the leptonic forward-backward asymmetry (with respect to
806 the direction of production) give information on the couplings of the Z' . At higher pp energies, the discovery
807 reach increases to 12 TeV for 33 TeV and 30 TeV for 100 TeV.

808 Lepton colliders are sensitive to new vector bosons that interfere with the s -channel virtual photon and Z in
809 two-fermion production $\ell^+\ell^- \rightarrow f\bar{f}$. The reach for discovery of a sequential Z' at the ILC at 500 GeV is also
810 about 7 TeV and scales proportional to the center of mass energy for higher energy colliders. Measurements
811 of the Z' signal with two possible beam polarizations and with individual lepton and quark final states gives
812 a large amount of information toward the identification of the quantum numbers of the boson.

813 1.8.2 Supersymmetry

814 Searches for supersymmetry (SUSY) encompass a wide range of strategies aimed at different particles of the
815 SUSY spectrum.

816 The most generic searches assume that supersymmetric partners of the SM particle carry a conserved
817 quantum number, called R-parity. If the lightest supersymmetric particle is neutral, it will typically be
818 weakly interacting and will not be observed in a collider detector. Events are then characterized as containing
819 several hadronic jets, associated with decay to the lightest particle plus missing transverse momentum. The
820 results of these analyses are parametrized by limits on the gluino mass and on a squark mass, assumed
821 common to all squark flavors. Current LHC results exclude such events up to gluino masses of 1.0 TeV and,
822 independently, squark masses of 1.3 TeV. For the future stages of the LHC, we expect to be able to discover
823 such events up to gluino masses of 1.9 TeV and squark masses of 2.3 TeV with 300 fb^{-1} , and to 2.3 TeV
824 and 2.7 TeV with 3000 fb^{-1} . This reflects more than a factor 2 in increased search power at the 300 fb^{-1}

825 stage, and another 20% with the additional factor 10 in luminosity. The gluino discovery reach increases to
 826 4.8 TeV for a 33 TeV pp collider and to 10.2 TeV for a 100 TeV collider.

827 The dependence of search reach on luminosity deserves comment. Away from kinematic limits for a given
 828 collider energy, parton-parton luminosity functions scale such that increasing the parton-parton center of
 829 mass energy by a factor 2 decreases the luminosity by a factor 10. This rule, which implies that a factor of 10
 830 in luminosity increases search reach by a factor of 2 in mass, must break down at masses near the kinematic
 831 limit. At the 14 TeV LHC, the reach increase falls off from the canonical factor of 2 for pair-produced
 832 particles with masses well above 1 TeV.

833 These considerations are important for models in which the first signal of SUSY would not be given by the
 834 generic search just described, but would require a more specialized analysis. Examples of such models are
 835 those in which mass gaps in the SUSY spectrum are relatively small (“compressed spectrum”), so that hard
 836 jets are not emitted in particle decays, and models in which only the partners of top quarks, or perhaps,
 837 only color-singlet supersymmetric particles are produced at accessible energies. Such models can be highly
 838 motivated. A compressed spectrum is needed in many models of supersymmetric dark matter particles to
 839 allow “coannihilation” to produce the correct dark matter density [40]. Models with only the top squarks
 840 and Higgsinos light satisfy the naturalness ideas of Sec. 1.2.2 in a minimal way. Reach estimates for top
 841 squark pair production were given above in Sec. 1.7.4.

842 Models in which SUSY discovery is more difficult at the 8 TeV LHC thus benefit more from the increase in
 843 luminosity to the HL-LHC. Models for which the first signal of SUSY would be the partners of W and Z
 844 bosons can be searched for at the 14 TeV LHC, with discovery expected up to masses of about 500 GeV.
 845 The factor of 10 luminosity increase to HL-LHC increases the reach by a factor of 2 in the analyses [41, 42],
 846 consistent with the argument given above.




Figure 1-7. PLACEHOLDER NP report fig. 1-20, holes at 300 and 3000

847 Another way to look at this issue is shown in Fig. 1-7. The figures shows a survey of a large number of SUSY
 848 models [43] plotted in the plane of gluino mass versus lightest superparticle mass. The color-coding gives
 849 the fraction of models excluded by LHC searches, at 300 fb^{-1} on the left, and at 3000 fb^{-1} on the right.
 850 The general shift of the boundary by about 30% is accompanied by a removal of exceptions to the right of
 851 this boundary.

852 One more exception should be noted. The supersymmetric partners of the Higgsino automatically have small
853 mass splitting of a few GeV. The direct pair-production of these particles through electroweak interactions
854 is essentially invisible at the LHC, except through searches for initial-state radiation plus invisible particles,
855 described in Sec. 1.8.4. In a scenario in which these are the lightest supersymmetric particles, the ability
856 of lepton colliders to be sensitive to very small energy depositions in decay would be crucial to observe and
857 study these particles. Studies of Higgsino pair production at the ILC as described in [44].

858 1.8.3 Long lived particles

859 The searches we have described so far assume that all new particles decay promptly. However, there are
860 many models that give exceptions to this. ATLAS and CMS have carried out dedicated searches for tracks
861 associated with long-lived massive particles and for particles decaying in the detector, perhaps out of time
862 with the bunch crossings. Current limits are stronger than those in searches for promptly decaying particles.
863 For example, ATLAS places limits of 310 GeV on a tau slepton, 600 GeV on a top squark, and 985 GeV
864 on a gluino. Should such long-lived particles exist, the LHC detectors trap a sample of them for detailed
865 studies of their decay modes and lifetimes.

866 1.8.4 Dark matter

867 The signature of jets plus missing transverse momentum discussed above for SUSY applies to a broader class
868 of theories. Any model in which the new TeV spectroscopy gives rise to a dark matter candidate particle
869 requires that this particle be kept stable over the age of the universe. Heavier states carrying QCD color
870 will decay to this particle, producing events of this same characteristic type. If the partners of quarks are
871 fermionic rather than bosonic, the production cross section will be higher. The generic SUSY search just
872 described can then be interpreted as a robust search for models predicting a TeV spectroscopy and dark
873 matter.

874 It is possible that the heavier states of the TeV spectrum are out of reach kinematically. Then the discovery
875 of dark matter would require the observation of direct pair-production of the essentially invisible dark matter
876 particles. This can be done using the fact that the production of particles a hard scattering process typically
877 also produces gluons or photons radiated from the initial-state particles. If the particles mediating the pair-
878 production reaction are heavy enough, this initial-state radiation can be at higher momentum than, and,
879 thus, distinguishable from, the ordinary particle production in typical collisions.

880 Reach estimates for the discovery of dark matter pair-production has been studied systematically in [45],
881 using an effective operator formalism to describe the coupling of dark matter to SM particles. This formalism
882 also allows cross sections measured at colliders to be related to rates for direct and indirect detection of dark
883 matter. Some results of this analysis, comparing limits on the dark matter-nucleon cross section from LHC
884 and higher energy pp colliders to limits from direct detection, are shown in Fig. 1-9. It is noteworthy that
885 a 100 TeV pp collider can place limits on the dark matter particle mass above 1 TeV, close to the unitarity
886 limit for thermal production of such a particle in the early universe.




Figure 1-8. TBA

887 1.8.5 The Message

888 The conclusions of the New Particles and Forces working group can be summarized as follows:

- 889 1. TeV mass particles are needed in essentially all models of new physics. The search for them is
890 imperative.
- 891 2. LHC and future colliders will give us impressive capabilities for this study. Future programs target
892 new physics at the all-important TeV scale, as can be seen in Fig. 1-8. *We need to get a version of this*
893 *plot that includes CLIC.*
- 894 3. This search is integrally connected to searches for dark matter.

895 Experiments on new particles and forces give information on the Particle Physics Questions # 1, 2, 3, 4, 5,
896 8, 9 listed in the Snowmass Summary [27].

897

898 1.9 Flavor Mixing and CP Violation at High Energy

899




Figure 1-9. NP report fig. 1-20, holes at 300 and 3000

900 The explanation of Higgs condensation may or may not give insight into the theory of flavor. It is a mystery
 901 why quarks and leptons have a hierarchical mass spectrum, why the weak interactions are not diagonal in
 902 flavor and violate CP. In the SM, these features are parametrized by the fermion-Higgs Yukawa couplings.
 903 The idea that that flavors are distinguished only through terms of the structure of the Yukawa couplings is
 904 called “Minimal Flavor Violation.”

905 Models of new physics at the TeV scale introduce a large number of new couplings. These in principle
 906 can be proportional to flavor-violating couplings with completely new structures. Such structures are not
 907 needed to build a model of Higgs condensation. In fact, many specific models—most notably, SUSY with
 908 gauge-mediated breaking [52]—are flavor-blind. It is possible also that some principle, analogous to the GIM
 909 mechanism, requires that new flavor couplings are related to the Yukawa couplings. In these cases, there are
 910 no new flavor-changing effects beyond the SM arising from the TeV scale.

911 However, it is also possible that flavor couplings among new particles have a different pattern. Such couplings
 912 can generate rare flavor-changing weak decays. They can also affect the phenomenology of the new particles
 913 themselves, requiring new strategies for searches. In this section, we discuss examples of models of this type.
 914 There are many possibilities, only a few of which can be discussed here. A more complete catalog is given
 915 in the working group report [46].

1.9.1 SUSY with Flavor-Dependent Soft Masses

In Sec. 1.8, we discussed reach estimates for models of TeV spectroscopy that were either blind to flavor or singled out only the third generation. More general forms of the new particle spectrum are allowed. The most important difficulties with flavor observables come when squarks with the same gauge quantum numbers, *e.g.*, the partners of d_R and s_R , have different masses. But there is little difficulty in giving the partners of d_R and d_L smaller masses than those of the other squarks. This weakens the experimental limits on the lightest squark mass. In fact, it is possible, with other mechanisms to suppress flavor effects of new physics, to allow only the partner of c_R to be light. This has been explored in detail for SUSY models [47, 48]. The current limit on the charm squark in such models is about 300 GeV.

1.9.2 R-parity Violating SUSY

It is possible that the R-parity conservation law that should keep the lightest SUSY partner stable is violated by new interactions. These interactions necessarily have a complex structure in flavor. One possible form of the R-parity violating interaction is

$$\lambda_{ijk}^1 U_i D_j D_k, \quad (1.17)$$

where U_i , D_i are the right-handed quarks or their squark partners and $i = 1, 2, 3$ is the generation number. This violates baryon number, allowing a squark to decay to two antiquarks. The interaction must be antisymmetric in color, and this requires it also to be antisymmetric in the flavors of the two down-type quarks. Other possible R-parity violating interactions have the forms

$$\lambda_{ijk}^2 L_i L_j \bar{E}_k, \quad \lambda_{ia}^3 L_i H_a \quad (1.18)$$

where L_i , \bar{E}_i are left-handed leptons or antileptons or their slepton partners and H_a is a Higgs or Higgsino field. These interactions violate lepton number. The coefficients of these operators are usually taken to be small to avoid unwanted flavor-changing rare decays. In particular, either the operator (1.17) or the lepton number violating operators must be highly suppressed to avoid rapid proton decay.

In R-parity violating models with the operator (1.17) only, SUSY decay chains typically end with the lightest SUSY particle decaying to jets. These jets must have a nontrivial flavor structure, possibly with a b jet always included.

R-parity violation with the Higgs operator in (1.18) can produce neutrino masses through the ‘‘Type III seesaw’’: A neutrino converts to a Higgsino, which then converts back to an antineutrino, possibly of a different flavor. The difference between the quark and lepton flavor mixing patterns is explained by the statement that the mass matrices come from unrelated operators. In such models, the branching ratios of the Higgsino are related to the neutrino mixing angles, and this relation can be confirmed by direct measurement [49]. In more general seesaw models of neutrino mass, it is possible that the seesaw scale could be the TeV scale, setting up other relations between TeV mass neutral leptons and the neutrino mixing matrix [50].

1.9.3 Models with Electroweak Baryogenesis

To explain the asymmetry between the numbers of baryons and antibaryons in the universe, a new source of CP violation is needed beyond that of the CKM phase. One natural place to look for this is in an

951 extended Higgs sector. A second requirement is that the electroweak phase transition be first-order. Then
 952 the transition takes place through the growth of bubbles, with top quarks scattering from the bubbles
 953 because they are massless outside and massive inside. A CP phase in the Higgs sector makes this scattering
 954 asymmetric between matter and antimatter. Both criteria can be satisfied in models with multiple Higgs
 955 fields [51]. Measurements of the Higgs self-coupling and of CP violation in Higgs decays, described in
 956 Secs. 1.4.2 and 1.4.3 above, test models of this type.

957 1.9.4 The Message

958 The conclusions of the Flavor and CP working group can be summarized as follows:

- 959 1. TeV mass particles may or may not introduce couplings with new types of flavor violation. These
 960 possible new couplings affect the search methods for new particles, in many cases, requiring new
 961 strategies.
- 962 2. The search for new particles is integrally connected to searches for rare flavor changing decays.

963 Experiments on flavor associated with new particles give information on the Particle Physics Questions #
 964 1, 2, 3, 4, 5, 8, 9 listed in the Snowmass Summary [27].

965 1.10 Scientific Cases for Future Colliders

966 In the previous sections presented the physics opportunities for the next steps in the Energy Frontier in terms
 967 of individual particles and research areas under study. It is also interesting to assemble these topics in terms
 968 of the experimental program that each accelerator in the list given in Sec. 1.3.2 will provide. Integrating
 969 over the topics in this way, we see that many of these proposed accelerators have very substantial physics
 970 programs that will explore the TeV energy scale across a broad range of measurements.

971 In this section, we present the cases for the various accelerators as if each accelerator stood on its own,
 972 with no further physics discoveries between now and the time that it begins operation. However, one should
 973 always keep in mind the possibility of discoveries would open up the study of physics beyond the SM. We
 974 have argued already that the likelihood of the discovery of new particle is very high even for the coming
 975 runs of the LHC at 14 TeV up to 300 fb⁻¹. Such a discovery would need to be followed up by further
 976 exploration that would benefit from accelerators with complementary capabilities or higher energy. This
 977 might, in the end, be the most important benefit of building the accelerators that come later in the timeline
 978 below. We will expand on this idea in Section 1.11.

979 1.10.1 LHC in This Decade: 300 fb⁻¹

980 First of all, we emphasize the many opportunities that will be provided by the coming run of the LHC
 981 at 14 TeV. Operation of the LHC at 14 TeV up through the 300 fb⁻¹ with the Phase 1 upgrades offers a
 982 tremendous increase in the power of new particle searches, close to a factor 2 in mass in most channels.
 983 Many of these searches, for example, the search for the gluino almost to 2 TeV and the search for vectorlike
 984 top partners above 1 TeV, access ranges of the masses that are strongly motivated in models of Higgs
 985 condensation.

986 This impressive capability is only one aspect of a broad program that will be carried out at the LHC over
 987 the next ten years. Its features include:

- 988 1. Clarification of Higgs boson couplings, mass, spin, CP to the 10% level.
- 989 2. First direct measurement of top-Higgs boson couplings
- 990 3. Precision W boson mass measurement below 10 MeV.
- 991 4. First measurements of VV scattering.
- 992 5. Theoretically and experimentally precise top quark mass to 600 MeV.
- 993 6. Measurement of top quark couplings to gluons, Z and W bosons, and photons with a precision
 994 potentially sensitive to new physics—a factor 2-5 better than today.
- 995 7. Search for top squarks and top partners and $t\bar{t}$ resonances predicted in models of composite top, Higgs.
- 996 8. New generation of PDFs with improved gluon and antiquark distributions.
- 997 9. Precision study of electroweak cross sections in pp collisions, including a new photon PDF.
- 998 10. Extension by a factor 2 in the sensitivity to new particles: SUSY, Z , top partners—key ingredients for
 999 models of the Higgs potential—and the widest range of possible TeV-mass particles.
- 1000 11. Deep ISR-based searches for dark matter particles.

1001 1.10.2 High-Luminosity LHC: 3 ab^{-1}

1002 The second high luminosity running of the LHC is referred to as the “Phase 2” upgrade period with
 1003 instantaneous luminosities of $5 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$. This running with 3,000 fb^{-1} of accumulated data truly
 1004 inaugurates the high-precision electroweak era at LHC with few percent precision for most Higgs boson
 1005 couplings as well as the 5 MeV threshold in M_W mass determination.

- 1006 1. The precision era in Higgs boson couplings begins, with sensitivities to 2-10%, and 1% for the ratio of
 1007 $\gamma\gamma$ and ZZ couplings.
- 1008 2. Measurement of rare Higgs boson decays, $\mu^+\mu^-$ and $Z\gamma$, with 100 M Higgs bosons.
- 1009 3. First measurement of the Higgs boson self-coupling.
- 1010 4. Deep searches for extended Higgs bosons.
- 1011 5. Precision W mass to ± 5 MeV.
- 1012 6. Precise measurements of VV scattering with access to Higgs sector resonances.
- 1013 7. . Precision top mass to ± 500 MeV.
- 1014 8. Deep study of rare, flavor-changing, top couplings with 10G tops.
- 1015 9. Search for top squarks and partners in models of composite top quarks and Higgs bosons in the expected
 1016 range of masses.
- 1017 10. Further improvement of q , g , and γ PDFs to higher x and Q^2 .

- 1018 11. A 20-40% increase in mass reach for generic new particle searches which can be as much as a 1 TeV
1019 step in mass reach.
- 1020 12. Extension by a factor of 2 in the mass reach for particles produced by the electroweak interactions.
- 1021 13. Any discovery at LHC — or in dark matter or flavor searches — can be followed up.

1022 1.10.3 ILC, up to 500 GeV

1023 The ILC would run at 250 GeV, 350 GeV, and 500 GeV, in a program that could begin as early as the
1024 second half of the next decade. It will study the properties of the Higgs boson, the top quark, and possibly
1025 also newly discovered particles, in very fine detail.

- 1026 1. Tagged Higgs boson study in $e^+e^- \rightarrow Zh$: model-independent Higgs boson width and branching ratio
1027 measurements, direct study of invisible and exotic Higgs boson decays
- 1028 2. Model-independent Higgs boson couplings with percent-level accuracy, and great statistical and sys-
1029 tematic sensitivity to BSM theories.
- 1030 3. Higgs boson CP studies in fermionic channels (e.g., $\tau^+\tau^-$).
- 1031 4. Giga-Z program for EW precision, W boson mass to 4 MeV and beyond.
- 1032 5. Improvement of triple vector boson couplings by a factor 10, to an accuracy below expectations for
1033 models with Higgs sector resonances.
- 1034 6. Theoretically and experimentally precise top quark mass to 100 MeV.
- 1035 7. Sub-% measurement of top couplings to gamma and Z, with accuracy well below expectations in models
1036 of composite top quarks and Higgs bosons.
- 1037 8. Search for rare top couplings in $e^+e^- \rightarrow t\bar{c}, t\bar{u}$.
- 1038 9. Improvement of α_s from the Giga-Z program.
- 1039 10. No-footnotes search capability for new particles in LHC blind spots – Higgsino, stealth stop, compressed
1040 spectra, WIMP dark matter.

1041 1.10.4 ILC at 1 TeV

1042 In the farther future, the extension of ILC to 1 TeV will access additional Higgs boson reactions for precision
1043 study and, possibly, also reach new particle thresholds.

- 1044 1. Precision measurement of the Higgs boson coupling to top, to 2% accuracy.
- 1045 2. Measurement of the Higgs boson self-coupling, to 13% accuracy
- 1046 3. Model-independent search for extended Higgs boson states to 500 GeV.
- 1047 4. Improvement in precision of triple gauge boson couplings by a factor 4 over 500 GeV results.
- 1048 5. Model-independent search for new particles with coupling to γ or Z to 500 GeV

- 1049 6. Search for Z using $e^+e^- \rightarrow f\bar{f}$ to about 5 TeV, a reach comparable to LHC for similar models. Multiple
1050 observables for diagnostics of the Z couplings.
- 1051 7. Any discovery of new particles dictates a lepton collider program: search for electroweak partners, 1%
1052 precision mass measurement, the complete decay profile, model-independent measurement of cross
1053 sections, branching ratios and couplings with polarization observables, search for flavor and CP-
1054 violating interactions

1055 1.10.5 CLIC: 350 GeV, 1 TeV, 3 TeV

1056 Extremely high energies in e^+e^- collisions will likely require technologies beyond that envisioned for the ILC.
1057 CLIC is a likely candidate facility which that would probe Higgs boson self-couplings and exotic scattering
1058 of both standard model particles and any new particles found or hinted at in earlier machines.

- 1059 1. Precision Higgs boson coupling to top, 2% accuracy.
- 1060 2. Higgs boson self-coupling, 10%.
- 1061 3. Model-independent search for extended Higgs **boson** states to 1500 GeV.
- 1062 4. Improvement in precision of triple gauge boson couplings by a factor 4 over 500 GeV results.
- 1063 5. Precise measurement of VV scattering, sensitive to Higgs **boson** sector resonances.
- 1064 6. Model-independent search for new particles with coupling to gamma or Z to 1500 GeV: the expected
1065 range of masses for electroweakinos and WIMPs.
- 1066 7. Search for Z using $e^+e^- \rightarrow f\bar{f}$ above 10 TeV.
- 1067 8. Any discovery of new particles dictates a lepton collider. program as with the 1 TeV ILC.

1068 1.10.6 Muon Collider: 125 GeV, 350 GeV, 1.5 TeV, 3 TeV

1069 A muon collider holds promise as a technique for reaching very high energies in lepton-lepton collisions and
1070 for s -channel production of the Higgs boson and possible additional Higgs states. Studies of the muon collider
1071 are not yet mature, particularly in designing a detector that can overcome the background from decays of
1072 the muons circulating in the ring. However, promising first results were reported at Snowmass.

- 1073 1. Similar capabilities to e^+e^- colliders described above.
- 1074 2. Ability to produce the Higgs boson, and possible heavy Higgs bosons, as s -channel resonances. This
1075 allows a sub-MeV Higgs boson mass measurement and a direct Higgs boson width measurement.

1076 1.10.7 Photon Collider

1077 Another technique for producing Higgs bosons in the s -channel is to convert an electron collider to a photon
1078 collider by backscattering laser light from the electron beams. This allows resonance studies at 80% of the
1079 electron center-of-mass energy.

- 1080 1. Production of Higgs or extended Higgs bosons as s -channel resonances, offering percent-level accuracy
 1081 in $\gamma\gamma$ coupling.
- 1082 2. Ability to study CP mixture and violation in the Higgs sector using polarized photon beams.

1083 1.10.8 TLEP, Circular e^+e^-

1084 An e^+e^- collider in a very large tunnel offers the possibility of very large integrated luminosity samples at
 1085 250 GeV and below, especially if multiple detectors can be used simultaneously.

- 1086 1. Possibility of up to 10 times higher luminosity than linear e^+e^- colliders at 250 GeV. Higgs boson
 1087 couplings measurements might still be statistics-limited at this level.
- 1088 2. Precision electroweak programs that could improve on ILC by a factor 4 in $\sin^2\theta_w$, a factor 4 in M_W ,
 1089 and a factor 10 in M_Z .
- 1090 3. Search for rare top couplings in $e^+e^- \rightarrow t\bar{c}, t\bar{u}$ at 250 GeV.
- 1091 4. Possible improvement in α_s by a factor 5 over Giga- Z , to 0.1% precision.

1092 1.10.9 A pp Collider: 100 TeV

1093 One of the ideas at Snowmass that gained momentum through the week was renewed interest in a Very
 1094 Large Hadron Collider. Reinvigorating R&D in a VLHC was a clear recommendation of the New Particles
 1095 and Forces Group and the conveners.

- 1096 1. High rates for double Higgs boson production; measurement of triple Higgs boson couplings to 8%.
- 1097 2. Deep searches, beyond 1 TeV, for extended Higgs boson states.
- 1098 3. Dramatically improved sensitivity to vector boson scattering and multiple vector boson production.
- 1099 4. Searches for top squarks and top partners and resonances in the multi-TeV region.
- 1100 5. Increased search reach over LHC, proportional to the energy increase, for all varieties of new particles
 1101 (if increasingly high luminosity is available). Stringent constraints on naturalness.
- 1102 6. Ability to search for electroweak WIMPs (e.g. Higgsino, wino), possibly covering the full allowed mass
 1103 range.
- 1104 7. Any discovery at LHC — or in dark matter or flavor searches — can be followed up by measurement
 1105 of subdominant decay processes, search for higher mass partners. Both luminosity and energy are
 1106 relevant.

1107 1.11 Discovery stories

1108 Another way to integrate over the physics topics presented in Sec. 1.4–1.9 is to consider the consequences of
 1109 a discovery of new physics at the LHC later in this decade. We have emphasized, first, that the presence of

1110 new particles at the TeV scale is necessary to build a physics explanation of electroweak symmetry breaking,
1111 and, second, that the coming run of the LHC, up to 300 fb^{-1} , will improve the depth of searches for new
1112 particles by more than a factor of 2. The conclusion from these statements is that the discovery of new
1113 physics at the LHC is likely. This means that we should have a plan for following up this discovery and
1114 exploring its implications. This program of course depends on the nature of the particle discovered, so a full
1115 analysis would be presented as a large number of case studies. We give two examples here for illustration.
1116 Further examples of these “discovery stories” are presented in the working group reports.

1117 1.11.1 Well-Tempered SUSY

1118 The New Particles and Forces working group [39] considered in some detail the consequences of a particular
1119 SUSY model that could be discovered at the LHC with 300 fb^{-1} . This particular model had a gluino at
1120 1.9 TeV , squarks ranging in mass from 1.3 TeV to 2.6 TeV , and bino and Higgsino states near 200 GeV . The
1121 bino and Higgsino were assumed to mix so that the lightest supersymmetric particle (LSP) would be a dark
1122 matter particle with the correct thermally generated cosmic density. This dark matter scenario is called the
1123 “well-tempered neutralino” [53].

1124 The LHC at 300 fb^{-1} would observe a robust jets plus missing transverse momentum signal. The signal
1125 would be dominated by the decay of the lighter squarks to a quark jet plus the unobserved LSP. The mass
1126 difference could be measured from kinematic distributions. Assuming that the LSP was light, this would
1127 also give an estimate of the squark mass. With the measured cross section, this would favor SUSY over
1128 models with fermionic partners.

1129 The HL-LHC would produce some of the heavier squarks and the gluino. Detailed kinematic measurements
1130 would identify at least one more mass scale in the spectrum and give further evidence for the SUSY
1131 hypothesis. The direct production of electroweak states would not be observed at the LHC, because these
1132 states have a compressed spectrum.

1133 A lepton collider with center of mass energy of 500 GeV would be able to pair-produce the Higgsinos and
1134 observe their decays to the LSP. Measurement of the polarized cross sections would give information about
1135 the quantum numbers of the electroweak states. It would also give an indirect determination of the mass of
1136 the electron-type slepton (750 GeV in this model) to 10 GeV . Using this information, it would be possible
1137 to evaluate the LSP annihilation cross section and show that it was consistent with that required for a dark
1138 matter particle.

1139 Experiments at higher-energy colliders would be needed to discover the heaviest sleptons (at 3.3 TeV) and
1140 squarks (at 2.6 TeV). Eventually, the complete SUSY spectrum would be determined, and the data on the
1141 mass spectrum could be used to deduce the pattern of SUSY breaking.

1142 1.11.2 $t\bar{t}$ Resonance

1143 An alternative scenario might be based on a Randall-Sundrum model with top and Higgs compositeness.
1144 The first evidence of this model would be the discovery of a resonance in pp collisions that decayed to $t\bar{t}$.
1145 Such a resonance at 3 TeV would be discovered at the LHC with 300 fb^{-1} . Study of kinematic distributions
1146 of the $t\bar{t}$ final state would reveal that the top quarks were highly polarized, a prediction of this model.

1147 The HL-LHC would discover an electroweak singlet top quark partner, and, possibly also, a doublet of
1148 quarks with vectorlike coupling to the electroweak interactions. It is possible that very accurate studies
1149 of the $t\bar{t}$ spectrum would also reveal the presence of a color-singlet resonance, somewhat below 3 TeV. Its
1150 higher-statistics study of the TeV resonance might reveal at decay to $t\bar{c}$ with branching ratio 10^{-3} .

1151 A lepton collider at 500 GeV would observe a significant 3% enhancement of the right-handed top quark
1152 coupling to the Z boson. The Higgs boson coupling to $\gamma\gamma$ might be enhanced at the 2% level by the
1153 radiative correction from the top quark partners. This would be discovered by combining the high statistics
1154 measurement of $BR(\gamma\gamma)/BR(ZZ^*)$ from the HL-LHC with the precise measurement of the Higgs coupling
1155 to Z at a lepton collider.

1156 These measurements would give a tantalizing first glimpse of the structure of the underlying composite Higgs
1157 model. Experiments at higher energy colliders capable of producing resonances up to 20 TeV in mass would
1158 be needed to explore the full structure of the spectrum of states.

1159 1.12 Conclusions

1160 In this report, we have described the future program of research at high-energy colliders and summarized
1161 the efforts of six Snowmass 2013 Energy Frontier working groups. The detailed conclusions that we have
1162 reviewed come back repeated to a set of four points that deserve special emphasis.

1163 **1.** The question of new particles at the TeV mass scale is a central issues in particle physics. We argued that
1164 new particles must exist to provide a physics explanation for the properties of the Higgs field. We need to
1165 know their nature, and their implications for the laws of physics at very short distances. What is the nature
1166 of the spectrum of new particles at the TeV mass scale? We argued that new particles must exist to provide
1167 a physics explanation for the properties of the Higgs field. We have given many examples in which these
1168 particles address other major questions of particle physics, including the questions of flavor, dark matter,
1169 and the unification of forces.

1170 **2.** The central capability of high-energy collider experiments is to produce massive elementary particle directly.
1171 In Energy Frontier experiments, we observe the W and Z bosons, the top quark, and the Higgs boson as real
1172 particles whose production and decay we can study in detail. The same will be true for any new particles
1173 that we can create. This is a unique, direct, and powerful method to learn about the laws of physics.

1174 **3.** There are three essential aspects in the exploration of the physics of the TeV mass scale.

1175 1. We must study the properties of the Higgs boson in as much detail as possible.

1176 2. We must search for the imprint of the TeV mass particles on the heaviest SM particles, the W and Z
1177 bosons, and the top quark.

1178 3. We must search for the direct production of the new particles at high energies.

1179 **4.** This program can be realized at accelerators now envisioned to operate in the coming years.

1180 1. We have emphasized the great opportunity that is being provided by the coming operation of the LHC
1181 at 14 TeV. The next stage of the LHC will double the range of searches for new particles and give
1182 similar leaps in capability for other probes of TeV physics.

- 1183 2. We have projected quantitatively what will be achieved in running the LHC at high luminosity, to
1184 ultimately acquire 3000 fb^{-1} of data per experiment. For some physics topics, the gain is incremental,
1185 but for others, in particular, the precision study of Higgs couplings and the search for new particles
1186 with only electroweak interactions, we move to a qualitatively new level. Looked at in total, this is a
1187 highly motivated physics program.
- 1188 3. We have listed many essential contributions to the exploration of the TeV scale that can be provided
1189 by lepton colliders. These include precision studies of the Higgs boson, the W and Z bosons, and the
1190 top quark, capable in all cases of discovering percent-level corrections predicted as the effects of TeV
1191 particles. The construction and operation of the ILC in Japan will realize these goals.
- 1192 4. We have emphasized that the quest to understand the TeV scale will not be finished with the results of
1193 accelerators of the next generation. We believe it likely that the discovery of new particles at the next
1194 stage of collider physics will open a definite path for exploration to still higher energies. That journey
1195 begins with renewed effort to bring advanced accelerators capable of higher energies to reality.

1196 We emphasized in our introduction that the discovery of the Higgs boson changes everything. This discovery
1197 points to potentially profound modifications of the laws of physics at energies relatively close to those we
1198 now access at accelerators. The quest for new phenomena and the insights they will provide has just begun.

1199 ACKNOWLEDGEMENTS

1200 The Energy Frontier conveners are profoundly grateful to the 26 dedicated conveners of our working groups:
1201 Kaustubh Agashe, Marina Artuso, John Campbell, Sally Dawson, Robin Erbacher, Cecilia Gerber, Yuri
1202 Gershtein, Andrei Griksan, Kenichi Hatakeyama, Joey Huston, Ashutosh Kotwal, Heather Logan, Markus
1203 Luty, Kirill Melnikov, Meenakshi Narain, Michele Papucci, Frank Petriello, Soeren Prell, Jianming Qian,
1204 Reinhard Schwienhorst, Chris Tully, Rick Van Kooten, Doreen Wackerroth, Lian-Tao Wang, and Daniel
1205 Whiteson.

1206 We are also grateful to the many scientists who assisted us in various ways through the process. We have
1207 benefited from those who have given us technical advice, from those who maintained our contact with the
1208 major collaborations, and from those who took time to help us with individual tasks.

1209 Our technical team was: Jeff Berryhill, Sergei Chekanov, Tom LeCompte, Sanjay Padhi, Eric Prebys, Tor
1210 Raubenheimer, and Eric Torrence. We also thank Markus Klute and Mark Palmer from the Capabilities
1211 group.

1212 Our advisors from the major experiments were: for ATLAS: Ashutosh Kotwal; for CMS: Jim Olsen; for
1213 LHCb: Sheldon Stone; for ILD: Graham Wilson; for SiD: Andy White; from CLIC: Mark Thomson; from
1214 the Muon Collider: Ron Lipton; and for the VLHC: Dmitri Denisov.

1215 Finally, we thank our fellow Snowmass 2013 conveners who asked “tough questions” and answered them.

References

- [1] See, for example, S. Schael *et al.* [ALEPH and DELPHI and L3 and OPAL and SLD and LEP Electroweak Working Group and SLD Electroweak Group and SLD Heavy Flavour Group Collaborations], Phys. Rept. **427**, 257 (2006) [hep-ex/0509008].
- [2] Examples of these mechanics are found, for supersymmetry, in L. E. Ibanez and G. G. Ross, Phys. Lett. B **110**, 215 (1982), L. Alvarez-Gaume, J. Polchinski and M. B. Wise, Nucl. Phys. B **221**, 495 (1983); for Little Higgs modes, in N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, JHEP **0207**, 034 (2002) [hep-ph/0206021]; for extra-dimensional models Y. Hosotani, S. Noda and K. Takenaga, Phys. Lett. B **607**, 276 (2005) [hep-ph/0410193].
- [3] Examples are found, for supersymmetry, in S. Weinberg, Phys. Rev. Lett. **50**, 387 (1983), H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983) [Erratum-ibid. **103**, 099905 (2009)]; for extra-dimensional models, in H. -C. Cheng, J. L. Feng and K. T. Matchev, Phys. Rev. Lett. **89**, 211301 (2002) [hep-ph/0207125].
- [4] Examples are found, for supersymmetry, in S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. D **24**, 1681 (1981); for extra-dimensional models, in K. Agashe, R. Contino and R. Sundrum, Phys. Rev. Lett. **95**, 171804 (2005) [hep-ph/0502222].
- [5] The following discussion oversimplifies an extensive and sophisticated literature on naturalness in supersymmetry. See, *e.g.*, R. Barbieri and G. F. Giudice, Nucl. Phys. B **306**, 63 (1988), B. de Carlos and J. A. Casas, Phys. Lett. B **309**, 320 (1993) [hep-ph/9303291]; J. L. Feng, K. T. Matchev and T. Moroi, Phys. Rev. Lett. **84**, 2322 (2000) [hep-ph/9908309]; M. Papucci, J. T. Ruderman and A. Weiler, JHEP **1209**, 035 (2012) [arXiv:1110.6926 [hep-ph]]; H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev and X. Tata, arXiv:1306.2926 [hep-ph] (Snowmass paper).
- [6] W. Barletta, *et al.*, Snowmass paper.
- [7] ref. on the LHC future
- [8] ref. on High-E LHC at 33 TeV
- [9] ref. on VLHC 100 TeV at CERN
- [10] ref. on VLHC 100 TeV
- [11] J. Brau, P. Grannis, M. Harrison, M. Peskin, M. Ross and H. Weerts, arXiv:1304.2586 [physics.acc-ph], Snowmass paper.
- [12] T. Behnke, *et al.*, arXiv:1306.6327 [physics.acc-ph], arXiv:1306.6352 [hep-ph], arXiv:1306.6353 [physics.acc-ph], arXiv:1306.6328 [physics.acc-ph], arXiv:1306.6329 [physics.ins-det].
- [13] ILC Higgs report? Snowmass paper.
- [14] D. Dannheim, P. Lebrun, L. Linssen, D. Schulte and S. Stapnes, arXiv:1305.5766 [physics.acc-ph]; H. Abramowicz *et al.* [CLIC Detector and Physics Study Collaboration], arXiv:1307.5288 [hep-ex], Snowmass paper.
- [15] M. Aicheler, *et al.*, CERN-2012-007.
- [16] J-P. Delahaye, C. Ankenbrandt, A. Bogacz, S. Brice, A. Bross, D. Denisov, E. Eichten and P. Huber *et al.*, arXiv:1308.0494 [physics.acc-ph].

- 1253 [17] Y. Alexahin, C. M. Ankenbrandt, D. B. Cline, A. Conway, M. A. Cummings, V. Di Benedetto, E. Eichten
1254 and C. Gatto *et al.*, arXiv:1308.2143 [hep-ph].
- 1255 [18] M. Koratzinos, *et al.*, arXiv:1305.6498 [physics.acc-ph]; M. Koratzinos, A. P. Blondel, R. Aleksan,
1256 P. Janot, F. Zimmermann, J. R. Ellis and M. Zanetti, arXiv:1306.5981 [physics.acc-ph], Snowmass
1257 paper.
- 1258 [19] S. A. Bogacz, J. Ellis, L. Lusito, D. Schulte, T. Takahashi, M. Velasco, M. Zanetti and F. Zimmermann,
1259 arXiv:1208.2827 [physics.acc-ph].
- 1260 [20] W. Chou, G. Mourou, N. Solyak, T. Tajima and M. Velasco, arXiv:1305.5202 [physics.acc-ph], Snowmass
1261 paper.
- 1262 [21] J. L. Abelleira Fernandez, C. Adolphsen, P. Adzic, A. N. Akay, H. Aksakal, J. L. Albacete, B. Allanach
1263 and S. Alekhin *et al.*, arXiv:1211.4831 [hep-ex].
- 1264 [22] S. Dawson, *et al.*, Snowmass paper.
- 1265 [23] H. E. Haber, in *From the Planck Scale to the Weak Scale (SUSY 94)*, P. Nath, T. Taylor, and S.
1266 Pokorski, eds. (World Scientific, 1995) [hep-ph/9501320].
- 1267 [24] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuszi and M. Spira, Eur. Phys. J. C **71**, 1753 (2011)
1268 [arXiv:1107.5909 [hep-ph]].
- 1269 [25] lattice writeup for Snowmass?
- 1270 [26] CMS Collaboration, Physics Analysis Summary CMS-PAS-HIG-13-005 (2013).
- 1271 [27] M. Bardeen, *et al.*, Snowmass paper.
- 1272 [28] A. Kotwal, *et al.*, Snowmass paper.
- 1273 [29] J. Campbell, *et al.*, Snowmass paper.
- 1274 [30] J. Beringer, *et al.* (Particle Data Group), Phys. Rev. **D86**, 010001 (2012).
- 1275 [31] W. -K. Tung, Acta Phys. Polon. B **33**, 2933 (2002) [hep-ph/0206114].
- 1276 [32] C. Anastasiou, L. J. Dixon, K. Melnikov and F. Petriello, Phys. Rev. Lett. **91**, 182002 (2003) [hep-
1277 ph/0306192].
- 1278 [33] M. Czakon, P. Fiedler and A. Mitov, Phys. Rev. Lett. **110**, 252004 (2013) [arXiv:1303.6254 [hep-ph]].
- 1279 [34] R. Aaij *et al.* [LHCb Collaboration], JHEP **1206**, 058 (2012) [arXiv:1204.1620 [hep-ex]].
- 1280 [35] K. Agashe, *et al.*, Snowmass paper.
- 1281 [36] G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, JHEP
1282 **1208**, 098 (2012) [arXiv:1205.6497 [hep-ph]].
- 1283 [37] S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **73**, 2494 (2013) [arXiv:1304.5783 [hep-ex]].
- 1284 [38] A. Abdesselam, *et al.*, Eur. Phys. J. C **71**, 1661 (2011) [arXiv:1012.5412 [hep-ph]].
- 1285 [39] Y. Gershtein, *et al.*, Snowmass paper.
- 1286 [40] K. Griest and D. Seckel, Phys. Rev. D **43**, 3191 (1991).

- 1287 [41] ATLAS Collaboration, arXiv:1307.7292 [hep-ex], Snowmass paper.
- 1288 [42] H. Baer, V. Barger, A. Lessa and X. Tata, arXiv:1306.5343 [hep-ph], Snowmass paper.
- 1289 [43] M. Cahill-Rowley, J. L. Hewett, A. Ismail and T. G. Rizzo, arXiv:1307.8444 [hep-ph], Snowmass paper.
- 1290 [44] M. Berggren, F. Brummer, J. List, G. Moortgat-Pick, T. Robens, K. Rolbiecki and H. Sert,
1291 arXiv:1307.3566 [hep-ph].
- 1292 [45] N. Zhou, D. Berge, T. M. P. Tait, L. Wang and D. Whiteson, arXiv:1307.5327 [hep-ex], Snowmass
1293 paper.
- 1294 [46] M. Artuso, *et al.*, Snowmass paper.
- 1295 [47] G. D. Kribs, E. Poppitz and N. Weiner, Phys. Rev. D **78**, 055010 (2008) [arXiv:0712.2039 [hep-ph]].
- 1296 [48] Y. Nomura, M. Papucci and D. Stolarski, Phys. Rev. D **77**, 075006 (2008) [arXiv:0712.2074 [hep-ph]].
- 1297 [49] J. List and B. Vormwald, arXiv:1307.4074 [hep-ex].
- 1298 [50] See, for example, P. Fileviez Perez, T. Han and T. Li, Phys. Rev. D **80**, 073015 (2009) [arXiv:0907.4186
1299 [hep-ph]].
- 1300 [51] D. E. Morrissey and M. J. Ramsey-Musolf, New J. Phys. **14**, 125003 (2012) [arXiv:1206.2942 [hep-ph]].
- 1301 [52] M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D **53**, 2658 (1996) [hep-ph/9507378].
- 1302 [53] N. Arkani-Hamed, A. Delgado and G. F. Giudice, Nucl. Phys. B **741**, 108 (2006) [hep-ph/0601041].