
Energy Frontier

1

2 [Tags, e.g. (Q#1), refer to items on the Big Question list.]

3

1.1 Energy Frontier

4 Experiments at the “Energy Frontier” make use of high-energy accelerators to produce and study heavy
5 elementary particles and to search for new ones. The Energy Frontier includes experiments at the Large
6 Hadron Collider (LHC) at CERN and those at future colliding-beam accelerators proposed for lepton-lepton
7 and proton-proton collisions.8 The first run of the LHC has closed a nearly half-century old chapter in the story of elementary particle
9 physics. We have discovered a most unusual new particle with properties very similar to those expected of
10 the Higgs boson. The appearance of this particle—and further confirmation of its identity—ends one era
11 and opens another. On one hand, the Standard Model of particle physics is complete. We know all of the
12 particles in this model and how they interact with one another and we have at least a basic idea of their
13 properties. On the other hand, we also know that the Standard Model is *incomplete* in important ways. It
14 challenges us to uncover the physics behind its apparently *ad hoc* structure. We are certain that a host of
15 observed anomalous phenomena and set of confusing conceptual questions have explanations that require
16 new physics outside the Standard Model.17 The LHC and the CMS, ATLAS, and LHCb detectors have brought to bear impressive capabilities for
18 exploring for the answers to these new questions. The LHC accelerator is expected to dramatically increase
19 its ability to deliver beams in the period between now and 2030, increasing its energy by almost a factor
20 of two and its integrated luminosity by a factor of 100. The detectors will improve their ability to collect
21 enormous data sets and to discriminate the properties of events with increasing precision. Around the world,
22 other new accelerators are being considered that will give us additional power in understanding the heaviest
23 particles of the Standard Model and exploring for new ones. In this report, and in the detailed working
24 group reports, we trace out the programs of these accelerators and present their most important goals.25 Our successful theory of weak interactions is based on the idea of an underlying symmetry that is spon-
26 taneously broken. The symmetry of the theory of weak interactions dictates the couplings of the quarks
27 and leptons to the W and Z bosons in a structure that has been confirmed by high-precision experiments.
28 However, this symmetry forbids the quarks, leptons, and vector bosons from having mass. The inclusion of
29 mass within the Standard Model requires a condensate of fields that fills the universe. The Higgs field—the
30 field of which the Higgs particle is the quantum—is the simplest realization of this idea.31 The discovery of the Higgs particle gives us an answer to the “how” question of mass generation. But
32 it does not answer the “why” questions: We have no understanding of the mechanism behind the Higgs
33 field condensation. We have no understanding of the origin of the couplings of quarks and leptons to the
34 condensate. We have no idea how to compute the couplings of the Higgs to quarks and leptons nor the
35 potential energy that leads to its condensation. Finally, all of the Standard Model particles come in groups,

36 yet the Higgs boson stands alone with properties that are unlike those of any other particle. This by itself
 37 is a puzzle, especially since its characteristics lead to more questions than they serve to answer. If these
 38 properties of the Higgs field and its quantum have physics explanations, there must be new particles and
 39 forces, and their masses cannot be far above the mass scale of the Higgs boson itself.

40 *These puzzles imply that new particles with masses of the order of 1 TeV which resolve these questions will*
 41 *be found—and will be accessible to existing and planned accelerators. (Q#9).* The discovery of the Higgs
 42 particle and acceptance of the challenges that it poses obligate us to carry out a three-pronged program of
 43 research in colliding beam experiments:

- 44 1. First, we must study the Higgs boson itself in as much detail as possible, searching for signs of a larger
 45 Higgs sector structure and the influence of new heavy particles.
- 46 2. Second, we must search for the imprint of the Higgs boson and its possible partners on the couplings
 47 of the W and Z bosons and the top quark.
- 48 3. Finally, we must search for the direct production of new particles with TeV masses predicted by models
 49 of the Higgs field and its symmetry breaking.

50 The Energy Frontier study pointed to all three of these approaches as important motivations for further
 51 experiments at colliders. The results of the study confirmed that the existing LHC detectors and their
 52 planned upgrades, together with proposed precision lepton collider experiments, will be nimble enough and
 53 sensitive enough to carry this three-fold campaign forward into the next two decades.

54 The Energy Frontier study was organized into six working groups, each associated with a physics topic. Each
 55 working group was asked to evaluate the future program for its subject both from a high-level perspective
 56 and from the viewpoint of supplying motivation for experiments at a range of proposed accelerators. In the
 57 remainder of this section, we will present the conclusions of these reports, first by physics topic, then by
 58 facility.

59 We begin with the Higgs boson. The bosonic resonance at 125 GeV was discovered at the LHC only one year
 60 ago. Its properties are now known to about the 30% level and, up to this point, its properties are consistent
 61 with those of the Higgs boson of the minimal Standard Model. The couplings of this boson roughly scale
 62 with mass. The specific form of the coupling to the Z boson indicates that the particle has spin-parity 0^+
 63 and that the corresponding field has a nonzero vacuum expectation value.

64 However, we cannot be complacent about the identity and role of this particle (Q#1). On one hand, the
 65 idea that a single scalar field is solely responsible for the generation of all particle masses is just a guess
 66 and needs explicit verification. On the other hand, models with additional Higgs bosons and related new
 67 particles, and models in which the Higgs boson is composite, are hardly tested. Deviations from the minimal
 68 Higgs boson properties due to new particles with mass M are suppressed by a factor $(m_h/M)^2$, so to the
 69 extent that the LHC has set lower limits on the masses of new particles at many hundreds of GeV, we would
 70 not yet have expected to see the modifications to the Higgs properties caused by those particles.

71 An experimental program to probe the Higgs boson contains several elements. The first is to search for
 72 deviations from the minimal Standard Model expectation that the Higgs boson couples to each particle
 73 species according to its mass. Such deviations are expected in almost all models of new physics, with a
 74 characteristic pattern for each model. The High-Luminosity LHC is expected to measure these couplings
 75 with accuracies of several percent, varying from coupling to coupling. Lepton collider experiments have the
 76 potential to push these accuracies to the sub-percent level, allowing discovery of percent-level deviations.
 77 Such a program of precision measurement of Higgs couplings requires a parallel concerted effort in precision
 78 theory. It also requires improvement of our knowledge of crucial input parameters such as α_s and m_b , which

79 can be provided by lattice gauge theory computations. Collider experiments can also probe the nonlinear
 80 Higgs field self-coupling to the 10-20% level, thereby testing the critically important question of the shape
 81 of the Higgs potential.

82 Future experiments on the Higgs particle should also improve our knowledge of its mass and quantum
 83 numbers. The spin of the observed resonance should already be clear from LHC data in this decade. A
 84 more subtle question is whether this particle contains a small admixture of a CP-odd state, signaling CP
 85 violation in the Higgs sector (**Q#4**) and confirmation of at least one additional Higgs-like particle. We
 86 discuss probes for this effect at various colliders.

87 Finally, it is important to search directly for additional Higgs bosons (**Q#9**). The LHC can probe to masses
 88 of 1 TeV with model-dependent limits. Lepton colliders can probe more model-independently to masses close
 89 to half of their collision energy.

90 The study of W and Z bosons has two aspects, first, the extension of the program of precision electroweak
 91 measurements, and, second, the search for perturbations of the three- and four-vector boson couplings.

92 The minimal Standard Model makes precise predictions for the well-studied precision observables M_W and
 93 $\sin^2 \theta_w$ (**Q#8**). At the moment, the observed values are within 2σ of the predictions; the deviations are
 94 consistent with the effects of new particles in a range of new physics models. Higher accuracy in this program
 95 is clearly needed (**Q#9**). Future experiments will sharpen our knowledge of these quantities and perhaps
 96 expose inconsistency with the Standard Model. The LHC, especially in its high-luminosity phase, has the
 97 potential to reduce the error on the W mass to $\pm 5 \text{ MeV}/c^2$. This requires a factor 7 decrease in the current
 98 error due to parton distribution functions and is a challenge to QCD researchers. Lepton colliders can make
 99 further improvements, to an error of $\pm 2.5 \text{ MeV}/c^2$ with a dedicated measurement of the WW threshold.
 100 A linear collider with beam polarization running at the Z resonance to produce 10^9 Z bosons (Giga- Z) is
 101 expected to reduce the error on $\sin^2 \theta_w$ by a factor 10. Finally, a circular e^+e^- collider operating in a 100
 102 km tunnel can potentially push both errors down by another factor 4. All of these precision measurements
 103 and their interpretation push on the inflexible correlations among the Standard Model particles and their
 104 respective forces. Such precision measurements in precision electroweak observables could be discoveries
 105 should the tight constraints within the Standard Model begin to unravel.

106 The second theme of W and Z studies is the search for anomalous nonlinear couplings of the vector bosons
 107 (**Q#8**). Collider experiments at higher energy are sensitive to the three-gauge-boson couplings and for the
 108 first time, to four-gauge-boson interactions, that is, direct measurement of vector boson scattering. Lepton
 109 collider experiments have the potential to push current uncertainties on three-boson couplings down by an
 110 order of magnitude, into the region in which new physics effects are predicted in models in which the Higgs
 111 boson is composite. Both hadron and lepton colliders can access vector boson scattering, but the total center
 112 of mass energy available to in a scattering process is a crucial factor. The high-luminosity LHC is sensitive
 113 to vector boson or Higgs resonances with masses well above 1 TeV.

114 Quantum Chromodynamics (QCD) is well established as the correct theory of the strong interactions.
 115 Nevertheless, advances in QCD are needed to achieve the goals of future experiments, especially at hadron
 116 colliders. For experiments at hadron colliders, improved knowledge of the parton distribution functions is
 117 needed. This can be achieved with data expected from the LHC on the rapidity distribution of W , Z , and
 118 top quark production. In addition, precision cross section computations, to the NNLO level, are needed for
 119 many 2- and 3-particle production processes, especially those involving the Higgs boson. This will require
 120 advances in the theoretical art of QCD computation. Finally, it is important to push the error on the value
 121 of α_s below the 0.5% level. Lattice gauge theory seems to be a promising avenue for achieving this.

122 The top quark was discovered at the Fermilab Tevatron and studied there with samples of tens of thousands
 123 of $t\bar{t}$ pairs. At the LHC, we will study top quarks in samples of billions. At future lepton colliders, we will use

124 the electroweak couplings of top quarks as a production mode and probe these with polarization observables.
125 Both methods will transform our knowledge of this quark, whose properties are intimately connected to the
126 mysteries of flavor and mass generation. To this day, we are surprised at the high mass of this presumably
127 fundamental particle and its proximity to the value of the Higgs vacuum expectation value.

128 The top quark mass is not only an important puzzle in itself but also is an important input parameter for
129 particle physics (Q#2). The strongest demands on precision in the top quark mass come from the precision
130 electroweak program, where interpretation of a $5 \text{ MeV}/c^2$ error in m_W requires a $500 \text{ MeV}/c^2$ error on m_t .
131 This mass must be a theoretically well-defined quantity, convertible to a short-distance parameter such as
132 the \overline{MS} mass. There are strategies applicable at the LHC which allow the measurement of a well-defined
133 top quark mass to this $500 \text{ MeV}/c^2$ accuracy. At lepton colliders, measurement of the cross section at the
134 top quark pair production threshold gives the \overline{MS} mass to $100 \text{ MeV}/c^2$, as required for the more accurate
135 precision electroweak program available at these machines.

136 Top quark couplings will be studied at high accuracy both at hadron and at lepton colliders. New physics from
137 top quark or Higgs compositeness can create few-percent corrections to the gluon, photon, and, especially,
138 Z boson couplings. These effects can be observed as corrections to the pair-production cross sections from
139 the predictions of the Standard Model. The top-quark coupling capabilities of a lepton collider is especially
140 strong, with accuracies possible at the sub-percent level (Q#1). The billions of top quarks produced at the
141 high-luminosity LHC allow very deep studies of rare flavor-changing top decays, to a level that complements
142 searches at low energy for flavor-changing quark decays (Q#2).

143 Models of the Higgs potential and its symmetry breaking typically require new particles that are partners,
144 in some way, of the top quark (Q#1). The LHC, especially in its high-luminosity stage, will have the
145 capability for deep searches for supersymmetric partners of the top quark, heavy vector-like top quarks that
146 appear in models with Higgs compositeness, and heavy resonances that decay to $t\bar{t}$, which appear in models
147 with new space dimensions (Q#9).

148 High energy colliders can search more generally for new particles with a very broad range of properties
149 (Q#9). These particles are required — and required to have masses near the 1 TeV scale — in models of
150 electroweak symmetry breaking. Others among our questions also call for new particles accessible to high
151 energy colliders. A large class of models of dark matter put the dark matter particle as the lightest particle
152 of a TeV mass spectroscopy (Q#5). Grand unification requires new particles near the TeV scale, including
153 partners of known particles and perhaps also new vector bosons associated with enhanced gauge symmetry
154 (Q#8). CP violation in the Higgs boson sector is required in models that generate the matter-antimatter
155 asymmetry at the electroweak phase transition (Q#4). More generally, new particles can bring new sources
156 of flavor and CP violation that might be reflected in the discovery of new flavor-changing reactions at low
157 energy (Q#2).

158 The LHC has already, in only its first run, increased the reach and power of searches for new particles
159 over a broad scope. We expect that this power will increase dramatically in the next decade, as the LHC
160 experiments acquire 300 fb^{-1} of data at 14 TeV. This extension probes deeply into the region expected
161 for new particles masses in all classes of models of electroweak symmetry breaking — and so, any plan for
162 high energy physics in the longer term must include the possibility of new particles discovery in this period
163 and exploitation of this discovery at the facilities that will follow. The high luminosity stage of the LHC,
164 up to 3000 fb^{-1} , will provide a further very significant extension of the search region. This extension is
165 particularly powerful for states produced through electroweak interactions, for which a factor of 2 increase in
166 the mass reach is available in some case. Lepton colliders would bring new and complementary capabilities,
167 in particular, the ability to carry out model-independent searches for states such as dark matter candidate
168 particles whose signatures are especially difficult at hadron colliders (Q#5).

169 The physics opportunities described above are reflected as physics motivations for current and future high
170 energy colliders. Our study considered a wide range of proposed machines. The full report from the Energy
171 Frontier presents the cases for these machines in some detail.

172 We first point out the opportunity provided by the 14 TeV run of the LHC schedule for the next decade.
173 This will provide robust searches for new particles over a broad front, with great promise of the discovery of
174 the TeV particle spectrum motivated at the beginning of this section.

175 We find the case for the high-luminosity stage of the LHC compelling. This plan to deliver 3000 fb^{-1} ,
176 has been listed by the European Strategy Study as the highest priority accelerator project in Europe for
177 the 2020's. We find that it will provide a significant additional step in the search for new particles, and
178 that it will provide other important capabilities. The most important of these is the beginning of the era
179 of precision Higgs boson measurements, to few-percent precision. It should give the first observation of
180 the Higgs boson self-coupling. It will provide a program of precision measurement in the Standard Model
181 that will dramatically tighten our knowledge of the W boson and the top quark, giving sensitivity in these
182 dimensions to a variety of new physics models. We have already noted that the additional luminosity will
183 significantly enhance the capability of the LHC to search for new heavy particles.

184 We considered the scientific case for the International Linear Collider. This next-stage lepton collider has
185 recently completed its Technical Design Report and is judged at Snowmass to be ready for construction.
186 This facility is named as the highest priority by the Japanese high energy physics community. We find the
187 case for this machine to be strongly motivated. It will reach sub-percent accuracy in the study of the Higgs
188 boson, allowing discovery of percent-level effects in the Higgs couplings predicted in new physics models.
189 It will measure the Higgs width in a model-independent way and will give the capability to observe Higgs
190 coupling to dark matter and other invisible and exotic modes of Higgs decay. It will extend our knowledge
191 of the top quark and the W and Z well below the percent level, setting up a confrontation with models that
192 include Higgs boson and top quark composite structure.

193 Many other accelerator facilities have been proposed to the Snowmass study for construction over longer
194 time scales. These include higher energy linear colliders, circular e^+e^- colliders, muon colliders, and photon
195 colliders. We include a detailed discussion of the physics motivations for these facilities in our full report.
196 There was particular interest in a proton collider of energy 100 TeV, which would come close to the capability
197 of covering the full model space for models of "natural" electroweak symmetry breaking and WIMP dark
198 matter. Our study developed materials and resources to begin a more complete study of physics at such a
199 high energy collider. This study, and a parallel development of magnet technology for higher-energy proton
200 colliders, should be pursued over the next decade.

201 Previous surveys of the prospects for high energy accelerator experiments have spoken in terms of reducing
202 the space of parameters—couplings, mixings, masses—as if that were the goal. Now more than ever, the
203 momentum points not toward exclusion, but toward the discovery of new states. Many possible directions
204 are open and must be pursued.

205 The Higgs boson discovery changes everything. It redefines the research agenda for particle physics, giving
206 us a set of sharp questions that we cannot ignore. The study of this unusual particle to high precision,
207 together with high precision studies of the W , Z , and top quark and searches for new states provide us
208 with complementary routes into the mysterious physics of symmetry breaking. The current LHC detectors
209 and their planned upgrades are well suited to carry on this program. Future accelerators will bring new
210 capabilities to pursue it further.

211 High energy colliders provide manifest opportunities to discover new fundamental interactions of broad
212 consequence. We believe that the US must be a part of these programs of discovery.