

- 34 • The quantum numbers of the quarks and leptons under the Standard Model gauge symmetry $SU(3) \times$
35 $SU(2) \times U(1)$ strongly suggests that these symmetry groups are unified into a larger *grand unification*
36 group $SU(5)$ or $SO(10)$; however, our precision knowledge of the strengths of the gauge couplings is
37 inconsistent with this hypothesis.
- 38 • The Standard Model cannot account for neutrino masses without the addition of some new particles.
- 39 • Further, the pattern of weak interaction mixing among neutrinos is completely different from that
40 observed for quarks.
- 41 • The Standard Model does not include the force of gravity or the small but nonzero energy in empty
42 space that gives rise to *dark energy*.

43 In addition, there is a major theoretical puzzle with the Standard Model. If the Higgs boson is an elementary
44 scalar particle, its mass is sensitive to the masses of any heavier particle to which it couples. It appears to
45 require a cancellation of one part in 10^{32} to explain why the Higgs boson mass is smaller than the Planck
46 mass.

47 The discovery of the Higgs boson has changed our viewpoint in how we address these questions, for three
48 reasons.

49
50 First, the Higgs boson completes the particle spectrum of the Standard Model. We have now discovered all
51 of the Standard Model particles and have measured many of their properties. It is clear now exactly what
52 the model does *not* explain. We have entered a new era in which the verification of the Standard Model
53 takes second place to a search for new, unknown forces and interactions.

54 Second, one of the key mysteries concerns the Higgs boson itself. The Higgs boson was predicted as a direct
55 consequence of the simplest model of the generation of mass for quarks, leptons, and gauge bosons. For
56 a long time, many particle physicists have expressed discomfort with this model. Now the prediction has
57 become a reality. We have to grapple with it and understand why nature chooses a particle with these
58 properties to do its work.

59 Third, the Higgs boson itself gives us a new experimental approach. Within the Standard Model, all properties
60 of the Higgs boson are precisely predicted from the value of the Higgs mass. But, as soon as we step outside
61 the Standard Model, the properties of the Higgs boson are hardly constrained by theory. It is compelling to
62 tug on this particle until the Standard Model breaks. We need to apply to the Higgs boson the same scrutiny
63 that we have applied in previous decades to hadron structure, heavy quark system, the W and Z bosons,
64 and top quark. Each study was done at the Energy Frontier machines of its day. This fruitful experimental
65 approach has acquired a new, promising target.

66 For exploration of the unknown regions outside the Standard Model, we are encouraged that very powerful,
67 experimental tools will be put into play. In the next ten years, the LHC at CERN is expected to almost
68 double its energy and to increase the size of its total event sample from the current 20 fb^{-1} to 400 fb^{-1} .
69 This new capability will put to the test many models that predict new physics beyond the Standard Model
70 and address the unexplained phenomena listed earlier in this section. In the decade after that, the LHC
71 should increase its data set by a further factor of ten. Lepton colliders and higher energy hadron colliders
72 are now receiving serious consideration for construction. The mysteries associated with the Higgs boson call
73 for new particles and forces at the TeV energy scale or the attometer distance scale. We now have before us
74 capabilities for a thorough exploration of this region of masses and distances. This is a compelling program;
75 the purpose of this report is to describe it in detail.

76 The structure of this summary report is as follows: In Section 1.2, we present the arguments for new
77 fundamental interactions at the TeV energy scale and the experimental program at colliders that these

78 arguments motivate. Section 1.3, we describe the organization of the Energy Frontier study. In Sections 1.4–
 79 1.9, we review in a more specific way the physics issues of collider experiments at the TeV energy scale. We
 80 consider in turn the prospects for exploration of new physics through studies of the Higgs boson, the W and
 81 Z bosons, Quantum Chromodynamics (QCD), the top quark, and searches for and study of new particles.
 82 We present the questions that need to be answered and the methodologies to attack these questions. In
 83 Sections 1.10, we present the capabilities of current and proposed colliders in relation to these physics goals.
 84 Finally, in Section 1.11, we trace out the implications of two possible scenarios for the discovery of new
 85 physics at the LHC. This gives an orthogonal way of appreciating the contributions that might be made by
 86 proposed accelerators. Section 1.12 gives our conclusions.

87 1.2 Importance of the TeV Scale

88 We have listed a number of motivations for new fundamental interactions beyond the Standard Model (SM).
 89 Where will we find them?

90 Explanations for baryogenesis, higher unification, and dark energy span a bewildering range of mass and
 91 distance scale. However, many of the questions we have listed in the previous section relate specifically to
 92 the energy scale of hundreds to thousands of GeV that we are exploring today at the Large Hadron Collider.
 93 We consider it imperative to understand particles and forces at this “TeV scale” thoroughly, using all of the
 94 tools at our disposal. In this section, we will discuss the importance of this regime of energies and short
 95 distances.

96 There is a sharp boundary at which our well-founded knowledge of the fundamental elementary particle
 97 interactions runs out. This is related to two different faces that the SM presents, which stand on very different
 98 theoretical foundations. On one side are the Yang-Mills gauge interactions, on the other side, the interactions
 99 of the Higgs field. The Yang-Mills interactions of quarks, leptons, and vector bosons are tightly determined
 100 by their quantum numbers and the strength of the coupling constants of the $SU(3) \times SU(2) \times U(1)$ vector
 101 bosons. Precision tests of the SM confirm the structure of these interactions to impressive accuracy [1].
 102 There is little doubt that the SM is a correct representation of nature at the energies we have currently
 103 explored.

104 On the other hand, the interactions of the SM fermions with the Higgs field, and the dynamics of the
 105 Higgs field itself, are essentially unconstrained and conceptually even cumbersome. The SM Lagrangian is
 106 constructed by writing down the most general terms allowed by gauge symmetry and renormalizability. The
 107 resulting potential term contains much of what is perplexing about the SM:

$$V = \mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2 + \sum_{f, f'} [g_{\Phi f f'} \bar{f}_L f'_R \Phi + h.c.] . \quad (1.1)$$

108 Here Φ is the spin 0 doublet field (doublet) and λ , $g_{\Phi f f'}$, and μ are parameters. The first two terms give
 109 the potential energy of the Φ field. When $\mu^2 < 0$, the neutral $I_3 = -1/2$ component of Φ is singled out and
 110 shifted by $\Phi^0 \rightarrow (v + h)/\sqrt{2}$, where $v = \sqrt{-\mu^2/\lambda}$ is the vacuum expectation value. At the same time, we
 111 diagonalize the matrix $G_{\Phi f f'}$. The last term in brackets then becomes

$$V(f f) = \sum_f \left[\frac{g_{H f f} v}{\sqrt{2}} \bar{f}_L f_R + \frac{g_{H f f}}{\sqrt{2}} \bar{f}_L f_R h + h.c. \right] . \quad (1.2)$$

112 Every term in (1.1) and (1.2) is of critical importance and each presents special challenges to interpretation
 113 and measurement. The first term in (1.2) is the new mass term for fermions, $m_f = g_{H f f} v/\sqrt{2}$. The pattern

114 of the fermion masses is totally unconstrained and proportional to the coupling constant g_{Hff} . The Yukawa
115 fermion-Higgs couplings (the second term of (1.2) and the angles from the diagonalization) contain the origin
116 of mass and mixings among quarks and leptons and CP violation in the weak interactions. Many parameters
117 in this sector are well measured, but there is no theory that explains their origin and structure.

118 As to the couplings of the Higgs boson itself (the first two terms of 1.1), the picture given in the Standard
119 Model is just one choice among many possibilities. There may be additional Higgs bosons and additional
120 particles of other types forming a larger “Higgs sector,” we have almost no information about these particles
121 except that their masses are probably larger than the mass of the known Higgs boson at 125 GeV,

122 1.2.1 The mystery of Higgs field symmetry breaking

123 The questions raised at the end of the previous section are brought to a focus by a single underlying question:
124 In order for any SM quark, lepton, or vector boson to obtain mass, the Φ field must condense and the resultant
125 Higgs field fills the universe. Why does this happen?

126 The SM itself provides no help with this question. It states only that symmetry breaking occurs if $\mu^2 < 0$,
127 which, as a physics explanation, is completely empty. Potentials of the form of (1.1) appear in many
128 condensed matter systems, including superconductors, magnets, and binary alloys. In those systems, it is
129 possible to compute the parameters of the potential from the underlying details of atomic structure and
130 explain why $\mu^2 < 0$. For the SM, if there is an underlying dynamics, its form is unknown. Attempts to
131 compute μ^2 within the SM, even to determine its sign, give disastrous results. The answers for μ^2 depend
132 quadratically on the values of large, unknown mass scales, with competing contributions of opposite sign.

133 Models are known in which μ^2 can be computed. However, they are not simple extensions of the SM. The
134 barrier to such a model is that the quadratic dependence on unknown scale parameters at very high energy
135 must be removed. However, this dependence is a generic property of models with fundamental scalar fields,
136 associated with the fact that the radiative corrections to the scalar field mass are quadratically divergent.
137 Cures for this problem require that the Higgs particle is non-generic in some important way: Either it is a
138 composite particle or it is related by a symmetry to a fermion or a vector boson. Symmetries of these types
139 can be included consistently only by profound extension of the structure of space-time, to supersymmetry
140 in the fermion case or higher dimensions of space in the vector boson case.

141 It is remarkable that, in each of these classes of models, easily identified radiative corrections give contri-
142 butions to μ^2 with a negative sign, predicting the instability of the Higgs field to condensation [2]. In all
143 three cases, these contributions come from quantum corrections due to partners of the top quark that are
144 predicted by the new symmetries.

145 The idea that the condensation of the Higgs field has a definite mechanical explanation from quantum
146 physics thus has major implications. It requires a new set of particles at the TeV mass scale. The examples
147 above include exotic partners of the top quark that are likely to be produced at the Large Hadron Collider.
148 This TeV particle spectroscopy can also supply explanations for other issues that require physics beyond
149 the SM. TeV particle spectra typically contain a massive neutral particle that can be absolutely stable and
150 thus a candidate for the particle of dark matter [3]. New couplings among the TeV particles potentially
151 provide new sources of CP violation, offering mechanisms for creating the matter-antimatter asymmetry of
152 the universe. Corrections to the SM coupling constants from the new particles can correct the evolution of
153 the SM couplings, allowing the three SM gauge interactions to unify at very short distances [4].

154 Most importantly, if the explanation for Higgs condensation changes our view of the SM itself — by making
 155 SM particles composite or by enlarging the structure of space-time — these changes must be taken into
 156 account in any explanation of phenomena that occur at still smaller distances scales, including the generation
 157 of neutrino masses, generation of flavor mixing among quarks and leptons, and the unification of the particle
 158 physics interactions with gravity.

159 In short: mechanisms that shed light on the physics behind the otherwise mysterious potential in Eq. 1.1
 160 are needed to directly address the major experimental anomalies of Section 1.1!

161 1.2.2 Naturalness

162 So the major question is: Are there any arguments that suggest to us how high in energy must we probe to
 163 discover the particles that address the questions of physics beyond the SM? There is no crisp answer. We
 164 do have a hint from the slippery principle called “naturalness.”

165 Naturalness is the statement that new particles that generate the μ^2 term in the Higgs potential (1.1) must
 166 have masses at the scale of μ^2 itself,

$$\mu^2 \sim (100 \text{ GeV})^2 . \quad (1.3)$$

167 Taken most naively, naturalness implies that new particles associated with the Higgs potential should have
 168 been found in the 1990’s at the experiments at LEP and the Tevatron. Today, the LHC experiments have
 169 carried out much deeper searches for these particles. How much further must we go?

170 One approach to naturalness looks more critically at the radiative corrections to the μ^2 parameter in the
 171 SM. The first-order corrections due to the top quark, the W and Z bosons, and the Higgs boson itself are

$$\delta\mu^2 = -\frac{3g_{Htt}^2}{8\pi^2}\Lambda^2 + \frac{3\alpha_w(3 + \tan^2\theta_w)}{4\pi}\Lambda^2 + \frac{\lambda}{8\pi^2}\Lambda^2 , \quad (1.4)$$

172 where g_{Htt} is the same Yukawa coupling as in (1.2), α_w and λ are the couplings of these particles, and θ_w
 173 is the weak mixing angle. All three terms are divergent, proportional to Λ^2 , where Λ is a mass scale at
 174 which the SM is replaced by a more complete underlying theory. Contributions from new particles add to
 175 (or subtract from) this expression. To give a well-defined result for μ^2 , they must cancel the dependence on
 176 Λ . If we allow the new contributions to cancel the SM ones over many decimal places, Λ can be arbitrarily
 177 high. However, this might be considered “unnatural.” If we assume that at most one significant figure is
 178 cancelled, we obtain interesting limits on top, W , and Higgs partners at roughly 2 TeV.

179 Another approach looks into the computation of μ^2 in specific models [5]. In supersymmetry (SUSY)
 180 models, the parameter called μ_{SUSY} — the Higgsino mass term — contributes to the Higgs parameter
 181 μ^2 at the tree level. Forbidding cancellations beyond one significant figure gives for the SUSY parameter
 182 $\mu_{SUSY} < 200 \text{ GeV}$. This is a strong upper bound on the mass of the supersymmetric partner of the Higgs
 183 boson, a particle that will be difficult to discover at the LHC. The supersymmetric partners of the top
 184 quark and the gluino contribute to the Higgs potential in one-loop and two-loop order, respectively. The
 185 corresponding naturalness bounds are

$$m(\tilde{t}) < 1 \text{ TeV} , \quad m(\tilde{g}) < 2 \text{ TeV} . \quad (1.5)$$

186 In Little Higgs models in which the Higgs boson is a composite Goldstone boson, the formula for the radiative
 187 correction to μ from a new fermionic partner T of the top quark has the form

$$\delta\mu^2 = C \frac{3\lambda_t^2}{8\pi^2} m(T)^2 , \quad (1.6)$$

188 where C is a model-dependent constant of order 1. This gives a bound

$$m(T) < 2 \text{ TeV} . \quad (1.7)$$

189 In all cases, we might have stronger cancellations in the expressions for μ^2 . Perhaps these cancellations might
 190 eventually find some physics explanation. However, each factor of 10 in mass above the bounds quoted
 191 requires cancellations of another *two* significant figures. Even such an imprecise criterion as naturalness
 192 probably limits top quark partners to lie below about 10 TeV.

193 However unsatisfactory these naturalness estimates might be, our interest in these estimates remains very
 194 strong. Higgs condensation is the mechanism that generates the whole spectrum of masses of the SM quarks,
 195 leptons, and vector bosons. Can it be just an accident? If not, there *must* be a spectrum of new particles
 196 at the TeV scale. Even if we cannot predict the value of this scale incisively, the importance of mass scale
 197 is clear. We must find these new states.

198 1.2.3 The mystery of dark matter

199 Independently of the naturalness argument, there is another, independent argument for new particles at the
 200 TeV mass scale. The Standard Model does not account for the dark matter which makes up 85% of the
 201 total matter content of the universe. Among the many explanations for dark matter, there is one class that
 202 is particularly compelling. This is the model of dark matter as composed of a neutral, weakly interacting,
 203 massive particle (WIMP) that was produced in the hot conditions of the early universe. The WIMP model
 204 of dark matter is discussed in full detail in the Cosmic Frontier report [6]. What is interesting here is
 205 one implication of the model: In order to obtain the observed density of dark matter, the energy scale
 206 of the interactions of dark matter must be close to 1 TeV. The TeV mass scale arises as a combination
 207 of astrophysical parameters with no obvious relation to the Higgs potential. Is this a coincidence, or a
 208 suggestion that models for the Higgs potential also solve the dark matter problem?

209 Theoretical models with WIMPs typically contain many particles with TeV scale masses. These particles
 210 share a common quantum number. They decay to the lightest particle carrying this quantum number, which
 211 is then stable for the lifetime of the universe. Whether or not this new spectrum of particles is connected to
 212 the Higgs problem, it is important to search for the production of those particles at colliders.

213 1.2.4 Summary

214 The ideas reviewed in this section predict a spectrum of new particles at the TeV mass scale. Those particles
 215 should be discoverable in experiments at the LHC and planned future accelerators. These experiments will
 216 provide the crucial tests of those ideas. Furthermore, if such particles are discovered, they can be studied in
 217 detail in collider experiments to determine their properties and to establish new fundamental laws of nature.

218 A research program in pursuit of new particles with TeV masses consists of three threads:

- 219 1. We must study the Higgs boson itself in as much detail as possible, searching for signs of a larger Higgs
 220 sector and the effects of new heavy particles.
- 221 2. We must search for the imprint of the Higgs boson and its possible partners on the couplings of the W
 222 and Z bosons and the top quark.

223 3. We must search directly for new particles with TeV masses that can address important problems in
224 fundamental physics.

225 To the extent that the naturalness or dark matter arguments above are a guide, all three approaches will be
226 accessible at high-energy collider experiments in the near future. In the next section, we will describe the
227 tools that we have available for that search.

228 1.3 Organization of the Energy Frontier study

229 In this section, we briefly describe how the Energy Frontier study was organized, in terms of topical working
230 groups and the landscape of proposed accelerators.

231 1.3.1 Working groups for the study of the Energy Frontier

232 We divided the study of the TeV energy scale thematically, in terms of probes of this scale using different
233 particles and interactions. The results summarized here constitute the efforts of hundreds of physicists who
234 worked through the winter and spring of 2013 within six working groups. The leaders of these groups are
235 co-authors of this report. The working groups were;

- 236 1. The Higgs Boson
- 237 2. Electroweak Interactions
- 238 3. Quantum Chromodynamics and the Strong Force
- 239 4. Understanding the Top Quark
- 240 5. The Path Beyond the Standard Model - New Particles, Forces, and Dimensions
- 241 6. Flavor Mixing and CP Violation at High Energy

242 Highlights of each group’s work are presented in this order in the following six sections. For each group, the
243 summary of results is followed by their “Message,” a quick summary of their conclusions. We follow with the
244 scientific cases to be made for each possible accelerator organized around each physics group’s conclusions
245 for that facility.

246 1.3.2 Accelerators for the Study of the Energy Frontier

247 In our discussion, specific estimates of the capabilities of the methods that we discussed will be made in the
248 context of proposed accelerator programs discussed at Snowmass. We provide here a brief orientation to
249 these programs. Energies refer to the center of mass energy of colliding beam experiments. For details on
250 the design and current status of these proposals, see the Capabilities Frontier working group report [7].

251 The baseline for our study is Large Hadron Collider (LHC), the pp collider now operating at CERN. The
252 most recent LHC schedule calls for 75-100 fb⁻¹ to be collected in a run starting in 2015 with, essentially,

253 the current detectors. Following Long Shutdown 2 in approximately 2019, the Phase 1 detector upgrades
 254 will be installed and running will resume at a projected instantaneous luminosity of $2 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$. This
 255 stage would produce another 300fb^{-1} of data. Then, in approximately 2023 the luminosity is expected to
 256 increase to $5 \times 10^{34} \text{cm}^{-2} \text{s}^{-1}$. In the Snowmass study, we compared the current results from the LHC, at
 257 7-8 TeV with an integrated luminosity of 20fb^{-1} , to future data samples at 14 TeV with 300fb^{-1} and with
 258 3000fb^{-1} . We refer to the latter program as the high-luminosity LHC or HL-LHC. The projected evolution
 259 of the LHC program is described in [8].

260 Our study considered higher energy pp colliders, with energy 33 TeV and 100 TeV. Unless it is indicated
 261 otherwise, the event sample assumed is 3000fb^{-1} . A high-energy upgrade of the LHC at 33 TeV (HE-LHC)
 262 is discussed in [9]. Colliders of 100 TeV energy are described in [10, 11]. In the following we will refer to
 263 such a collider generically as VLHC.

264 Our study considered e^+e^- linear colliders, both the International Linear Collider (ILC), covering the energy
 265 range 90 GeV–1000 GeV and the Compact Linear Collider (CLIC), covering the energy range 350 GeV–
 266 3000 GeV. The ILC is described in [12] and in its technical design report [13]. The TDR/CDR luminosity
 267 samples are 1000fb^{-1} at 1 TeV and scaling linearly with energy. Luminosity upgrades of the baseline ILC
 268 using strategies outlined in the TDR, to 2500fb^{-1} at 1 TeV and similar enhancements at other energies,
 269 with long running periods, are described in [14]. CLIC is described in [15] and in its Conceptual Design
 270 Report [16].

271 Our study considered $\mu^+\mu^-$ colliders operating over a range from 125 GeV to 3000 GeV. The luminosity
 272 samples assumed were similar to those for linear e^+e^- colliders. The technology of the muon collider is
 273 described in [17, 18].

274 Our study considered a circular e^+e^- collider in a large (80-100 km) tunnel. Accelerator parameters for such
 275 a machine are described in [19] in the context of one such proposal, TLEP, for a large tunnel near CERN. In
 276 principle, accelerator techniques invented for super-B-factors can produce very high luminosities, in excess
 277 of $10^{36}/\text{cm}^2\text{sec}$ at 90 GeV and $10^{35}/\text{cm}^2\text{sec}$ at 250 GeV, when summed over 4 detectors. However, there
 278 is as yet no complete accelerator design. The luminosities are lower, and/or the power consumption rises
 279 dramatically, at 350 GeV and above. In the following, we will refer to such a collider as TLEP (wherever it
 280 might be built). We will assume the above luminosities and operation with four detectors.

281 Two more types of accelerators received more limited attention from our study. Linear e^+e^- colliders
 282 can be converted to photon-photon colliders, with roughly 80% of the energy and similar luminosity, by
 283 backscattering laser light from the electron beams. Proposals for photon-photon colliders are described in
 284 [20, 21]. Colliding the LHC beam with an e^- or e^+ beam from a linear accelerator offers the opportunity of
 285 high energy ep collisions. This has been studied for a facility at CERN called LHeC, described in [22].

286

287 1.4 The Higgs Boson

288

289 We begin with the study of the Higgs boson itself. In this section, we will refer to the new boson with mass
 290 125 GeV as “the Higgs boson,” while recognizing that its properties could well be very different from the
 291 simplest expectations.

292 We have already emphasized that the study of the Higgs boson gives a completely new avenue along which to
 293 probe the physics of the TeV scale. The picture of the Higgs boson given by the SM is precise. All properties

of the Higgs boson can be computed now that the mass of the Higgs boson is known. And yet, this precise theory has no conceptual foundation. Current experiments exclude deviations from the SM at the 100% level, but surprises at the 30%, 10%, or 3% level are all possible in different highly plausible models. The nature of the Higgs boson is a central part of the mystery of TeV physics. New physics responsible for Higgs condensation must couple to the Higgs boson and affect its properties at some level.

Full details of the future program on the Higgs boson, and more precise statements of the uncertainty estimates given below, can be found in the Higgs Boson working group report [23].

1.4.1 Higgs boson couplings

The most direct question to ask about the new boson is that of whether it is in fact the sole source of mass for all quarks, leptons, and gauge bosons. For this to be true, the couplings of the Higgs boson to the various species of SM particles must follow a definite pattern. The couplings of the particle to fermions and vector bosons must be, from Eq. 1.2,

$$g_{Hf\bar{f},SM} = \frac{1}{\sqrt{2}} \frac{m_f}{v} , \quad g_{HVV,SM} = \frac{1}{2} \frac{m_V^2}{v^2} , \quad (1.8)$$

where v is the value of the SM Higgs condensate, equal to 246 GeV. (More properly, this is the leading-order prediction for a coupling defined to all orders with all three particles on shell.) These couplings have a simple pattern that should be tested for as many SM species as possible. In the following discussion, we define the scale factors

$$\kappa_A = g_{HA\bar{A}}/(SM) , \quad (1.9)$$

where (SM) denotes the SM prediction.

The Higgs boson also couples to pairs of vector bosons gg , $\gamma\gamma$, and γZ through loop diagrams. In the SM, these couplings are dominated by contributions from W boson and top quark loops. In more general theories, these couplings can also receive contributions from radiative corrections with new particles in loops. We will denote ratios of the on-shell couplings to the SM predictions by κ_g , κ_γ , $\kappa_{\gamma Z}$.

Corrections to the predictions (1.8) can appear at many levels. If there are multiple Higgs fields that mix into the observed boson, the κ_A will contain cosines of the mixing angles. These can be as large as the data permit. Radiative corrections due to loop effects of new particles are expected to be below the 10% level.

Corrections to the Higgs couplings are also affected by the Decoupling Theorem [24]: If all new particles have masses greater than M , we can integrate out these particles. The result is the SM, in which the properties of the Higgs boson are predicted precisely in terms of its mass. The corrections to the SM values are generated, in this way of constructing the formalism, by effective higher-dimension operators added to the SM. These corrections will then be at most of the order of m_h^2/M^2 . The Decoupling Theorem implies an apparently paradoxical but nevertheless important conclusion: In a model in which the Higgs sector is very complex but all new particles in it are heavier than 500 GeV, corrections to the Higgs boson properties are at most at the 5-10% level. We are likely to be in this situation, in which the picture of the Higgs boson may be very different from that in the SM but, since the other particles in the sector are heavy, it is difficult to conclude this except by precision measurement.

Typical sizes of Higgs boson coupling modifications are shown in Table 1-1. More details of these estimates are given in [23].

Model	κ_V	κ_b	κ_γ
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
Decoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$< 1.5\%$
Composite	$\sim -3\%$	$\sim -(3-9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$

Table 1-1. Generic size of Higgs coupling modifications from the Standard Model values when all new particles are $M \sim 1$ TeV and mixing angles satisfy precision electroweak fits.

Tests of the values of the Higgs couplings relative to the SM must take account of the theoretical uncertainty in the comparison to the SM predictions. A potentially observable quantity is the partial decay width $\Gamma(h \rightarrow A\bar{A})$, related to κ_A by

$$\kappa_A^2 = \Gamma(h \rightarrow A\bar{A}) / (SM) . \quad (1.10)$$

Currently, some of these quantities have large uncertainty in their evaluation in the SM. The uncertainty in the partial width $\Gamma(h \rightarrow b\bar{b})$, which accounts for more than half of the SM Higgs total width, is quoted as 6% [25]. A concerted program is required to bring the uncertainties in the SM predictions below 1%. This requires complete evaluation of the 2-loop electroweak corrections to the partial widths. It also requires improvement of the uncertainty in the crucial input parameters α_s , m_b , and m_c . Lattice gauge theory promises to reduce the errors on all three quantities to the required levels [26]. Further methods for improvement in our knowledge of α_s are discussed in Section 1.6.

There are only a few cases in which the partial widths $\Gamma(h \rightarrow A\bar{A})$ can be measured directly. More often, the Higgs decay partial widths are measured from the rates of reactions that involve the Higgs boson in an intermediate state. An example is the rate of $\gamma\gamma$ production through gg fusion at the LHC. The rate of this process is proportional to

$$\sigma(gg \rightarrow h) \cdot BR(h \rightarrow \gamma\gamma) \sim \frac{\Gamma(h \rightarrow gg)\Gamma(h \rightarrow \gamma\gamma)}{\Gamma_T(h)} , \quad (1.11)$$

where $\Gamma_T(h)$ is the total Higgs boson width. In terms of the κ_A quantities, the measured rates are proportional to

$$\sigma(A\bar{A} \rightarrow h)BR(h \rightarrow B\bar{B}) / (SM) = \frac{\kappa_A^2 \kappa_B^2}{\sum_C \kappa_C^2 BR_{SM}(h \rightarrow C\bar{C})} . \quad (1.12)$$

The SM prediction for the total width of the Higgs boson is 4 MeV, a value too small to be measured directly except at a muon collider where the Higgs boson can be produced as a resonance. At all other cases of hadron and lepton colliders, the total width must be determined by a fit to the collection of measured rates. Such fits entail some model-dependence to control the size of modes of Higgs decay that are not directly observed.

The report [23] compares the capabilities of the LHC and a variety of lepton colliders to extract the values of the Higgs boson couplings. At the LHC, the total number of Higgs bosons produced is very high, over 170 million per experiment for integrated luminosity of 3000 fb⁻¹. However, Higgs boson production at the LHC is accompanied by very high backgrounds. The extraction of couplings from cross sections is complicated by significant QCD uncertainties in the calculation of cross sections, currently about 12% for gluon fusion and 3% for vector boson fusion.

At electron colliders, the Higgs boson is produced in the relatively background-free interactions $\ell^+\ell^- \rightarrow Zh$ and $\ell^+\ell^- \rightarrow \nu\bar{\nu}h$ (vector boson fusion). The measurements of rates are dominated by statistical errors, but




Figure 1-1. Measurement precision on κ_γ at different facilities.




Figure 1-2. Measurement precision on κ_W at different facilities.




Figure 1-3. Measurement precision on κ_b at different facilities.




Figure 1-4. Measurement precision on κ_t at different facilities.

358 the statistics are limited. The Zh reaction offers tagged Higgs bosons, giving the possibility of observing
 359 decay modes not accessible at the LHC (such as decay to $c\bar{c}$), and invisible and exotic modes of Higgs
 360 decay. The total cross section for the two e^+e^- reactions are directly proportional to $\Gamma(h \rightarrow ZZ^*)$ and
 361 $\Gamma(h \rightarrow WW^*)$, respectively, without dependence on the $\Gamma_T(h)$. This allows lepton collider measurements
 362 to determine $\Gamma_T(h)$ and all individual partial decay widths by fitting of Higgs rates without any model
 363 assumptions.

364 Figures 1-1 through 1-4 show the comparison for a variety of accelerator programs of the projected
 365 uncertainty of measurement of the Higgs couplings. The first three figures show the uncertainties in the
 366 couplings to $\gamma\gamma$, WW , and $b\bar{b}$ from a 6-parameter fit appropriate to the analysis of LHC result. The fourth
 367 shows the projected error on the Higgs coupling to $t\bar{t}$ from experiments directly sensitive to this quantity.
 368 The facilities considered are the LHC at its current stage, after 300 fb^{-1} , and after 3000 fb^{-1} , the ILC
 369 to 500 GeV and up to 1000 GeV, the TLEP circular e^+e^- collider and the CLIC linear collider operating
 370 at 1400 and 3000 GeV. For LHC, the upper and lower estimates reflect a pessimistic scenario, in which
 371 systematic errors not evaluated from data do not improve, and an optimistic scenario in which theory errors
 372 are halved and all experimental systematic errors decrease as the square root of the integrated luminosity,
 373 For the ILC stages, the first error bar corresponds to the baseline event samples considered in the ILC TDR,
 374 while the second includes, more optimistically, a set of luminosity upgrades described in the TDR. Full
 375 details, and tables of the numerical results of the fit, can be found in [23]. The figures show, first, that the
 376 LHC, especially in its high luminosity phase, will measure Higgs couplings with impressively high precision.
 377 However, the discovery of perturbations of the Higgs boson couplings at the level shown in Table 1-1, at

378 3-5 σ significance, will require both the much lower level of systematic errors available at a lepton collider
 379 and very large event samples to reduce the statistical errors.

380 1.4.2 Higgs boson self-coupling

381 A particularly important coupling of the Higgs boson is the Higgs self-coupling, λ in (1.1), which determines
 382 the shape of the Higgs potential. In the SM, after Higgs condensation, there is a triple Higgs boson coupling
 383 proportional to $\sqrt{\lambda}$, given alternatively by

$$\lambda_{hhh} = \frac{3m_h^2}{v^2} . \quad (1.13)$$

384 This coupling can be extracted from the rate for double Higgs production, for example $pp \rightarrow hh + X$ or
 385 $e^+e^- \rightarrow \nu\bar{\nu}hh$.

386 Theoretical models with extended Higgs sectors or Higgs compositeness can predict deviations of the triple
 387 Higgs coupling of 20% relative to the SM expectation. These are larger effects than those expected in the
 388 Higgs couplings to fermions and vector bosons, but the measurement is also much more difficult. The cross
 389 sections at lepton colliders is at the fb level. At the LHC, the cross sections are larger, but the use of rare
 390 decay modes, including $h \rightarrow \gamma\gamma$, is considered to reduce background. Studies using only a few decay modes
 391 indicate a precision of 50% in the Higgs self-coupling measurement per detector; combining the results of
 392 the detectors, the first evidence of the Higgs self-coupling would be likely from the HL-LHC program. The
 393 projected uncertainties on the Higgs self-coupling are 13% in a long-term program at the ILC at 1 TeV or 10%
 394 for CLIC at 3 TeV. The double Higgs production cross section increases rapidly with energy. Measurements
 395 at a 100 TeV pp collider are estimated to reach an uncertainty of 8%.

396 1.4.3 Higgs boson spin and CP

397 A crucial test of the identification of the 125 GeV resonance with the Higgs boson is the measurement of
 398 its spin and parity. This issue is almost settled with the current data from the LHC. The fact that the
 399 resonance decays to $\gamma\gamma$ implies that it has integer spin and cannot have spin 1. The distribution of the four
 400 leptons in $h \rightarrow ZZ^*$ decays already strongly favors the 0^+ over the 0^- spin-parity hypothesis and excludes
 401 the simplest forms of spin 2 coupling [27]. This issue should be decided with the next LHC data set.

402 However, there is a more subtle issue associated with the Higgs boson CP. If there are multiple Higgs bosons
 403 and CP violation in the Higgs sector, the Higgs boson at 125 GeV can contain an admixture of CP scalar
 404 states. CP violation in the Higgs sector has major implications. Most importantly, it can provide the new
 405 source of CP violation outside the SM that allows the matter-antimatter asymmetry of the universe to be
 406 generated at the electroweak phase transition.

407 CP violation in the Higgs sector can be reflected both in production and decay of the Higgs boson. The
 408 most accurate tests are available in the study of the 4-lepton final state in $h \rightarrow ZZ^*$. CP-violating terms in
 409 this vertex can be masked by the large tree level decay amplitude proportional to the Higgs condensate v .
 410 However, measurements in vector boson fusion and in associated production are potentially accurate enough
 411 to overcome the loop suppression. Lepton colliders can search for CP violation in the decay $h \rightarrow \tau^+\tau^-$ and
 412 in the production process $\ell^+\ell^- \rightarrow t\bar{t}h$. The first process can reach 1% precision in the measurement of a
 413 CP-odd decay amplitude.

414 Photon-photon colliders, which produce the Higgs boson as a resonance, can use initial-state polarization
 415 to search for CP-violating terms in the Higgs boson coupling to $\gamma\gamma$, which has no tree-level contributions.
 416 Similarly, a muon collider can probe for CP-violating contributions to the Higgs boson coupling to $\mu^+\mu^-$ if
 417 the accelerator provides transverse beam polarization.

418 1.4.4 Higgs boson mass and width

419 The Higgs boson mass is currently known from the LHC experiments to better than 600 MeV. This accuracy
 420 is already sufficient for the uncertainty in the Higgs mass not to be significant in precision electroweak tests.
 421 The most important influence of a highly accurate Higgs mass within the SM comes in the evaluation of
 422 the predictions for the Higgs couplings to WW and ZZ , for which one boson must be off the mass shell. A
 423 100 MeV error in the Higgs mass corresponds to a 0.5% uncertainty in κ_W . We expect that the error in the
 424 Higgs mass can be decreased to 100 MeV and to 50 MeV, respectively, for the LHC programs with 300 fb^{-1}
 425 and 3000 fb^{-1} by using the $\gamma\gamma$, ZZ^* , and $\mu^+\mu^-$ modes in which the Higgs boson can be fully reconstructed.
 426 A lepton collider studying the Higgs boson in the Zh production mode would push this uncertainty down
 427 further, to about 35 MeV for linear colliders and 7 MeV for a very high luminosity program at a circular
 428 collider.

429 Predictions of the Higgs mass in models of new physics might provide further motivation for measuring
 430 the Higgs mass accurately. An example of such a model is the Minimal Supersymmetric Standard Model
 431 (MSSM). To evaluate the prediction to an accuracy of 100 MeV, however, the masses of the top squarks
 432 must be known, and the top quark mass must be known to 100 MeV.

433 We have noted already that lepton colliders offer the possibility of a model-independent determination of
 434 the Higgs boson total width. Because the couplings of the Higgs boson to ZZ and WW appear both in the
 435 expressions for measurable total cross sections and branching ratios, these couplings can be eliminated to
 436 evaluate the total width through the relations

$$\Gamma_T(h) \sim \sigma(\ell^+\ell^- \rightarrow Zh)/BR(h \rightarrow ZZ^*) \sim \sigma(\ell^+\ell^- \rightarrow \nu\bar{\nu}h, h \rightarrow b\bar{b})/BR(h \rightarrow WW^*)BR(h \rightarrow b\bar{b}) \quad (1.14)$$

437 This gives the Higgs boson width to 3% for a long-term program at the ILC and to 0.6% for a high luminosity
 438 program at a circular collider with multiple detectors. These uncertainties are reflected in the coupling
 439 uncertainties quoted in Section 1.4.1.

440 A muon collider would have the capability of observing the Higgs boson as a narrow resonance. For the
 441 projected beam energy resolution of 4×10^{-5} , the mass of the Higgs boson would be measured to 0.06 MeV
 442 and the width would be measured directly in the s -channel to a precision of 4% [18]

443 1.4.5 Searches for additional Higgs bosons

444 There are strong motivations for expecting the existence of additional Higgs particles. These motivations
 445 begin with the overall mysteries of the physics of Higgs condensation and the question of whether the Higgs
 446 boson is the only particle of the SM whose quantum numbers do not come in multiples. Beyond this, virtually
 447 all models of new physics to explain the Higgs potential contain additional Higgs doublet fields. These fields
 448 are required in supersymmetric models in order for Higgs fields give mass to both the up-type and the
 449 down-type quarks. In models with new space dimensions, additional Higgs fields arise as the Kaluza-Klein




Figure 1-5. 5σ discovery reach for a 300 GeV H decaying via $H \rightarrow ZZ \rightarrow 4\ell$, for the Type I and Type II two-Higgs-doublet models, shown in the left and right panels, respectively. The yellow regions show the 5σ reach in direct searches at the LHC. The successive larger regions correspond to the LHC at 300 fb^{-1} and 3000 fb^{-1} and a 33 TeV pp collider with 3000 fb^{-1} . The blue regions show the regions allowed at 95% CL by precision Higgs coupling measurements; the darker and lighter regions correspond to the LHC with 300 fb^{-1} and 3000 fb^{-1} [28].

450 excitations of the fundamental Higgs doublet. Each additional Higgs doublet gives rise to four new particles,
 451 CP-even and CP-odd neutral scalars H and A , and a charged pair H^\pm .

452 Often, extended Higgs bosons have enhanced couplings to heavy flavors, either to b and τ or to t depending
 453 on whether the extended Higgs parameter $\tan\beta$ is greater than or less than 1. This emphasizes searches
 454 based involve $b\bar{b}$ annihilation to Higgs resonances.

455 Currently, the LHC experiments based on heavy flavor signatures exclude additional Higgs bosons for masses
 456 as high as 1 TeV in restricted ranges of $\tan\beta$. The region of large $\tan\beta$ is surveyed by reactions such as
 457 $b\bar{b} \rightarrow H, A \rightarrow \tau^+\tau^-$, while the region of low $\tan\beta$ is surveyed by reactions such as $gg \rightarrow H, A \rightarrow t\bar{t}$,
 458 $gg \rightarrow A \rightarrow Zh$. A gap remains for intermediate values, roughly $2 < \tan\beta < 20$, which is closed only for
 459 extended Higgs boson masses up to 200 GeV. Future runs of the LHC, up to 3000 fb^{-1} , are expected to
 460 close this window up to about 500 GeV.

461 Additional Higgs particles typically have smaller couplings to WW and ZZ than the lightest Higgs boson.
 462 This coupling is proportional to the combination of mixing angles $\cos(\beta - \alpha)$, which is also constrained
 463 by measurement of the vector boson couplings to the known Higgs boson. The complementarity of these
 464 search strategies is illustrated with a search for additional Higgs bosons in a model with two Higgs doublets
 465 [28, 29, 30]. Fig. 1-5 shows the discovery reach for a 300 GeV Higgs boson H decaying via $H \rightarrow ZZ \rightarrow 4\ell$
 466 compared with allowed region from Higgs coupling measurements at various experimental facilities. The
 467 reach for a pseudoscalar A decaying via $A \rightarrow Zh \rightarrow \ell\ell b\bar{b}, \ell\ell\tau\tau$ is comparable [30]. The important parameters
 468 for this search are the Higgs mixing angle α and the ratio of the Higgs vacuum expectation values $\tan\beta$, and
 469 which Higgs bosons couple to which quarks and leptons (type I or II). The line $\cos(\beta - \alpha) = 0$ is the limit
 470 where the additional Higgs particles decouple from gauge bosons and the couplings of the 125 GeV Higgs
 471 boson are Standard Model-like. Even for Type II models (which include supersymmetry), where the Higgs
 472 coupling measurements are already very constraining, the direct Higgs search probes significant additional

473 parameter space. There is roughly a factor of 2 increase in the reach for large $\tan\beta$ in each step in going
474 from LHC14 to HL-LHC to a 33 TeV collider with 3000 fb^{-1} .

475 Lepton collider experiments can search for extended Higgs boson states through the reaction $\ell^+\ell^- \rightarrow HA$ up
476 to the kinematic limit, independently of the value of $\tan\beta$. The cross section depends only on the electroweak
477 quantum numbers of the extended Higgs particles. This covers the parameter space up to 500 GeV for ILC
478 and up to 1500 GeV for CLIC running at 3 TeV. Photon and muon colliders have the opportunity to discover
479 additional Higgs bosons as resonances up to the full center of mass energy of the machine.

480 1.4.6 The Message

481 The conclusions of the Higgs Boson working group can be summarized as follows:

- 482 1. Direct measurement of the Higgs boson is the key to understanding electroweak symmetry breaking.
483 The fact that the Higgs boson appears as a light, apparently fundamental, scalar particle needs
484 explanation. A research program focused on the Higgs couplings to fermions and vector bosons and
485 achieving a precision of a few percent or less is required to address these questions.
- 486 2. Full exploitation of the LHC is the path to few percent precision in the Higgs coupling and to a 50
487 MeV precision in the determination of the Higgs mass.
- 488 3. Full exploitation of a precision electron collider is the path to a model-independent measurement of the
489 Higgs boson width and a sub-percent measurement of the Higgs couplings. Such precision is necessary
490 to probe for new physics beyond the reach of LHC direct searches.

491 Experiments on Higgs bosons give information on the Particle Physics Questions # 1, 2, 4, 5, 8, 9, 10 listed
492 in the Snowmass Summary [31].

493 _____

494 1.5 Electroweak Interactions

495 _____

496 The precision electroweak experiments of the 1990's established the $SU(2) \times U(1)$ theory of electroweak
497 interactions at the sub-percent level of accuracy. But, they did more. All particle species with couplings to
498 the electroweak interactions eventually influence the properties of the weak interaction bosons W and Z . Very
499 precise measurements of the properties of these boson then have the potential to reveal new, undiscovered
500 particles. The experiments of the 1990's indicated the presence of a heavy top quark and a light Higgs boson
501 and estimated the masses at which these particles were eventually discovered. They disfavored a fourth
502 generation of quarks and leptons, now excluded by direct search at the LHC.

503 Increased precision in the properties of the weak interaction bosons could well turn up the first evidence of
504 the TeV spectrum of particles discussed in Section 1.2.1. There are still tensions evident in the data which
505 have been intriguing for years; for example, the current value of M_W from many experiments persists at
506 $1-2\sigma$ higher than the SM expectation. Experiments over the next decade will explore whether these could
507 become significant deviations requiring radiative corrections due to new particles.

508 In a description of possible new interactions in terms of effective operators, the electroweak precision
 509 observables probe only the first few terms. Experiments at higher energy probe additional operators by
 510 observing and constraining the nonlinear interactions of the W and Z . These operators can receive corrections
 511 from loop diagrams involving new TeV mass particles but, more strikingly, they can receive leading-order
 512 corrections if there is new strong dynamics or resonances in the Higgs sector.

513 We will review these topics in this section. Full details of the program, and more precise statements of the
 514 projected uncertainties described below, can be found in the Electroweak Interactions working group report
 515 [32].

516 1.5.1 Precision observables M_W and $\sin^2 \theta_w$

517 The current uncertainty in the W boson mass is 15 MeV, corresponding to an relative precision of 2×10^{-4} .
 518 It is remarkable that the most accurate determinations of M_W come from the hadron collider experiments
 519 CDF and DØ.

520 Precision measurement of M_W at hadron colliders is very challenging, but certain features of W production
 521 make it feasible to reach high accuracies. The directly measured transverse mass distribution is very sensitive
 522 to M_W , having a relatively sharp endpoint at the W mass. Likewise the p_T distributions of the leptons are
 523 also sensitive to the boson mass with different, but manageable systematic uncertainties. Enormous statistics
 524 will be available with very small contamination by background. The dominant errors come from the small
 525 corrections to these properties. Currently, the largest source of systematic error is the dependence of the
 526 acceptance on the rapidity of the produced W , requiring a correction that depends on quark and antiquark
 527 parton distribution functions (PDFs). Experimental uncertainties are at the same level as those due to PDFs
 528 and are expected to continue to decrease accordingly.

529 We see good prospects for improving this measurement at the LHC. The statistical component of the error
 530 will be negligible already with the current LHC data set. The error from PDFs doubles in going from the
 531 Tevatron to the LHC because proton-proton collisions give no valence antiquarks. However, we anticipate
 532 that this error will be decreased using new data on the vector boson rapidity and charge asymmetries. The
 533 issue of PDF improvement is discussed further in Section 1.6.2. The huge statistical precision will allow for
 534 control of calorimetric and tracking systematic uncertainties. In Table 1-2 we see that the PDF error in
 535 M_W can be brought down to ± 5 MeV with 300 fb^{-1} and to ± 3 MeV with 3000 fb^{-1} , leading to a final
 536 uncertainty in M_W of ± 5 MeV. In each stage, the experimental systematics are expected to keep pace. In
 537 order to reach this important level of precision, the PDF uncertainties must be pushed to a factor of 7 better
 538 precision that currently available.

539 Lepton colliders offer an opportunity to push the uncertainty in M_W down even further. The W mass was
 540 measured at LEP to ± 36 MeV from the kinematics of W^+W^- production. The uncertainty was dominated
 541 by statistical errors, with a substantial additional contribution from the modeling of hadronization. Both
 542 sources will benefit from the data set on W^+W^- , about 1000 times larger, that will be available at next-
 543 generation e^+e^- colliders such as ILC and TLEP. We estimate an error below ± 4 MeV from this method,
 544 and a similar error from independent measurements on single W production.

545 The ultimate W mass measurement would come from a dedicated energy scan of the W^+W^- threshold at
 546 160 GeV. Such a measurement could reach ± 2.5 MeV with the statistics available from the ILC and \pm
 547 1 MeV with the statistics available from TLEP. At this level, systematic errors become dominant. The
 548 program also requires a detailed precision theory of the W^+W^- threshold, using methods now applied to
 549 the $t\bar{t}$ threshold.

ΔM_W [MeV]	LHC		
\sqrt{s} [TeV]	8	14	14
\mathcal{L} [fb ⁻¹]	20	300	3000
PDF	10	5	3
QED rad.	4	3	2
$p_T(W)$ model	2	1	1
other systematics	10	5	3
W statistics	1	0.2	0
Total	15	8	5

Table 1-2. Current and target uncertainties in the measurement of M_W at the LHC.

550 The measurement of the value of $\sin^2 \theta_w$ associated with quark and lepton couplings to the Z resonance
 551 offers an orthogonal probe of the electroweak interactions. The current accuracy in $\sin^2 \theta_w$ is at the
 552 7×10^{-5} level of precision and is dominated by measurements from LEP and SLC. This level might be
 553 reached but, we expect, will not be surpassed at the LHC. Again, uncertainties in PDFs give the limiting
 554 systematic error. Measurements from the polarization-dependence of the Z cross section and from the b
 555 quark forward-backward asymmetry are discrepant by about 3σ , indicating an experimental question that
 556 should be resolved.

557 Future lepton colliders give an opportunity to improve the value of $\sin^2 \theta_w$. The ILC program includes a few
 558 months of running at the Z resonance to produce a data set of 10^9 Z 's, improving the statistics from LEP
 559 by a factor of 100 with highly polarized beams. The ILC detectors should also dramatically improve the
 560 capability for heavy flavor tagging. This ‘‘Giga- Z ’’ program should improve the uncertainty in $\sin^2 \theta_w$ by a
 561 factor 10. The program also would give new measurements of other Z pole observables sensitive to new TeV
 562 mass particles, most importantly, the fraction R_b of Z decays to $b\bar{b}$.

563 TLEP envisions a multi-year program at higher luminosity to collect 10^{12} events on the Z resonance.
 564 This potentially pushes the precision of electroweak measurements by another order of magnitude, though
 565 systematic contributions to the errors must still be understood. Among other factors, the Z mass must be
 566 measured more accurately than the current 2.5 MeV. This is possible at TLEP if transverse polarization can
 567 be achieved in single beam operation. The direct measurement of $\sin^2 \theta_w$ optimally requires longitudinal
 568 polarization in colliding beam mode; the feasibility of this at TLEP needs to be understood.

569 Loop effects from TeV mass particles can produce effects at the 10^{-4} level in both M_W and $\sin^2 \theta_w$, so
 570 the improved capabilities for precision electroweak measurements may point to new particle discovery or
 571 confirmation. Quantitative estimates for a number of models are given in [32].

572 1.5.2 Interactions of W and Z bosons

573 The interactions of W and Z bosons are studied at higher energies, through the measurement of vector
 574 boson pair production and multi-vector boson production. This study has already begun at LEP and the
 575 Tevatron, where parameters of the 3-gauge boson interactions were bounded within a few percent of their
 576 SM values.

577 Vector boson interactions are described in a unified way through the formalism of effective Lagrangians.
 578 One begins from the SM Lagrangian, in which the Yang-Mills vertices for γ , W , and Z appear as terms of
 579 dimension 4. One then adds higher dimension operators invariant under the $SU(2) \times U(1)$ gauge symmetry.
 580 A typical term involving an operator of dimension 6 is

$$\delta\mathcal{L} = \frac{c_W}{\Lambda^2} (D_\mu\Phi)^\dagger W_{\mu\nu} (D_\nu\Phi) , \quad (1.15)$$

581 where Φ is the Higgs doublet field and $W_{\mu\nu}$ is the W boson field strength. This operators contributes to
 582 the 3- and 4-vector boson vertices. Additional operators of dimension 8 can modify the 4-vector interactions
 583 independently of the 3-vector interactions. A typical term is

$$\delta\mathcal{L} = \frac{f_{T,0}}{\Lambda^4} \text{tr}(W_{\mu\nu})^2 \text{tr}(W_{\lambda\sigma})^2 . \quad (1.16)$$

584 In a weak-coupling theory such as the SM, the coefficients c_i and f_j are induced by loop diagrams and should
 585 be highly suppressed, by powers of $\alpha_w/4\pi \sim 10^{-3}$. However, in theories with strong interactions in the Higgs
 586 sector, the c_i and f_j coefficients could be of order 1, with the Λ parameters then interpreted as the masses
 587 of Higgs sector resonances. For example, the operator (1.16) would be induced by a scalar resonance in the
 588 Higgs sector.

589 The current bounds on triple gauge boson couplings imply that the Λ parameters associated with dimension
 590 6 operators are higher than about 600 GeV. High statistics measurements of the triple gauge bosons by
 591 observation of W^+W^- and ZZ production in e^+e^- reactions at 500 GeV are expected to be sensitive to
 592 deviations from the SM that are 10 times smaller, pushing the sensitivity to Λ almost to 2 TeV.

593 It is difficult for hadron colliders to have similar sensitivity to triple gauge couplings. One source of this
 594 difficulty is that the LHC experiments study diboson reactions at higher energies, where additional terms
 595 from higher dimension operators are important and so the extraction of the coefficients of dimension 6
 596 operators is model-dependent. But there is a compensatory advantage. Working at higher energy, the LHC
 597 will study W and Z bosons at energies where Higgs sector strong interactions can dramatically alter the
 598 amplitudes for vector boson scattering.

599 Some quantitative examples are presented in [32]. In examples studied there, the sensitivity to the coefficients
 600 f/Λ^4 of dimension 8 operators implies that, assuming $f = 1$, the LHC at 30 fb^{-1} would achieve exclusion of
 601 a Λ value greater than 1.5 TeV. The HL-LHC, with 3000 fb^{-1} , would roughly triple the significance of the
 602 effect, allowing 5 sigma discovery at 1.5 TeV or exclusion up to 1.8 TeV. If the origin of this new physics
 603 lies in strongly-coupled dynamics, the HL-LHC provides discovery-level sensitivity to virtual effects of new
 604 resonances with masses ($4\pi\Lambda$) up to 19 TeV. If interpreted as sensitivity to f , the HL-LHC is more sensitive
 605 by a factor of 2-3 compared to the LHC at 300 fb^{-1} .

606 Increasing the energy allowed for the diboson system dramatically increases the physics reach. For discovery
 607 of anomalous couplings, a 33 TeV pp collider would reach Λ values above 1.8 TeV, while a VLHC at 100 TeV
 608 would reach above 4.6 TeV. These values extend well into the region in which new Higgs sector dynamics
 609 would be expected in models of this type.

610 1.5.3 The Message

611 The conclusions of the Electroweak Interactions working group can be summarized as follows:

- 612 1. Precision measurements of the W and Z bosons has the potential to probe indirectly for new particles
 613 with TeV masses. This precision program is within the capabilities of LHC, linear e^+e^- colliders, and
 614 TLEP.
- 615 2. Measurement of vector boson interactions will probe for new dynamics in the Higgs sector. In such
 616 theories, we expect correlated signals in triple and quartic gauge couplings. The LHC and linear
 617 colliders will have sensitivity into the mass region above 1 TeV.

618 Experiments on electroweak interactions give information on the Particle Physics Questions # 1, 4, 8, 9
 619 listed in the Snowmass Summary [31].

620

621 1.6 Quantum Chromodynamics and the Strong Interaction

622

623 Every probe of particle physics at high energies eventually requires detailed knowledge of the strong inter-
 624 actions. Even in pure electroweak processes, the strong interactions affect the values of α and α_w through
 625 coupling constant renormalization. In the next decade, when most of our new knowledge about particle
 626 physics will come from hadron collider experiments, our understanding of the strong interactions will affect
 627 every aspect of the data, through the structure of the proton, through radiative corrections to initial- and
 628 final-state quarks and gluons and their transition to hadrons, and through the detailed physics that produces
 629 multi-jet events.

630 The strong interactions are known to be described by the Yang-Mills theory Quantum Chromodynamics
 631 (QCD). This is a theory that is weakly coupled at short distances and strongly coupled at large distances.
 632 Our understanding of QCD is imperfect. We have limited tools for the strongly coupled regime, and precision
 633 calculation in the weakly coupled regime is technically complex. Nevertheless, our knowledge of QCD has
 634 taken enormous strides since the previous Snowmass workshop a decade ago. In this section, we review the
 635 current state of our tools for QCD and indicate the opportunities for further progress. More details on all
 636 of the topics discussed here can be found in the working group report [33].

637 In our discussion of precision quantum field theory calculation, we will describe one-loop radiative corrections
 638 as next-to-leading order (NLO), and higher corrections as NNLO, *etc.*

639 1.6.1 α_s

640 The strength of the QCD coupling is determined by the value of the coupling constant α_s , usually expressed
 641 as its \overline{MS} value at the Z mass. The current Particle Data Group [34] value of this quantity has 0.6%
 642 uncertainty. We have pointed out in Section 1.4.1 that this is a limiting systematic error in the evaluation
 643 of the SM prediction for Higgs boson couplings.

644 There are three strategies to improve the estimate. The extraction of α_s from the Z boson width or the
 645 ratio of hadronic to leptonic Z decays is theoretically unambiguous but is limited by statistics. Currently,
 646 this extraction has a 4% uncertainty and does not improve the world average. The Giga- Z program at the
 647 ILC discussed in the previous section should improve this uncertainty to 0.4%. The very high luminosity Z
 648 program envisioned for TLEP could decrease this uncertainty further to 0.1%. We judge that higher-statistics
 649 measurements of e^+e^- event shapes are not competitive with these improvements.

650 Proposed improvements of PDFs from LHeC will lead to an improved value of α_s from the measurement
 651 of PDF evolution. The expected statistical error would be $\pm 0.2\%$ from LHeC alone and $\pm 0.1\%$ from the
 652 combination of LHeC and HERA. The theoretical systematic error for this method is not as well understood,
 653 but we estimate this at $\pm 0.5\%$ once one further order in QCD perturbation theory ($N^3\text{LO}$) is calculated.

654 The most accurate current values of α_s come from lattice gauge theory. Higher statistics lattice estimates
 655 and calculation of additional terms in lattice perturbation theory should decrease the current uncertainties
 656 over the next decade to 0.3% . These improvements will come together with improvements in the values of
 657 the quark masses, as discussed in 1.4.1.

658 1.6.2 Parton distribution functions

659 Our knowledge of the initial state in hadron-hadron collisions is encoded in the representation of the proton
 660 structure given by the parton distribution functions (PDFs). The provision of PDF distributions with
 661 uncertainties is an innovation of the past decade [35]. The uncertainties quoted have been continually
 662 improved through the addition of new data sets, especially from the Tevatron and HERA experiments.




Figure 1-6. Comparison of the partonic luminosities at the 14 TeV LHC, as a function of the parton center of mass energy, from the CT10, MSTW, and NNPDF2.3 NNLO PDF sets: Left: $q\bar{q}$ luminosity; Right: gg luminosity, from [33]

663 Still, there are gaps in our knowledge, especially in the components relevant to the study of physics beyond
 664 the SM. The leading PDF distributions disagree in their estimates of the gluon-gluon luminosity function at
 665 the mass of the Higgs boson; this accounts for an 8% systematic uncertainty in the extraction of the cross
 666 section for Higgs production. The step from the Tevatron to the LHC puts added weight on the antiquark
 667 distributions in the proton. And, all parton luminosities are poorly constrained by data for parton-parton
 668 invariant masses greater than about 500 GeV. This is clearly shown in Fig. 1-6.

669 We expect that these difficulties can be addressed using future data from the LHC. PDFs at the few-percent
 670 level of accuracy require theoretical calculations at the NNLO level. These are already available for the
 671 Drell-Yan process [36]. The NNLO computation of the total cross section for top quark pair production has
 672 recently been completed [37] and is now being extended to the rapidity distribution. NNLO calculations
 673 for 2-jet production are in progress. Over the next few years, these calculations will be used in conjunction
 674 with a very high-statistics data set on jet, top quark, and lepton pair production from the LHC. The LHCb
 675 experiment has an important role to play in the measurement of rapidity distributions at $y > 2.5$ [38].

676 Further improvements in PDFs can result from the program of the LHeC. The data expected will reduce
677 the error in the gluon luminosity to a few percent at the Higgs boson mass and to 5-10% in the multi-TeV
678 region.

679 1.6.3 Electroweak corrections to hadron collider processes

680 The quest for few-percent accuracy in predictions for hadron colliders brings new elements into play. In
681 particular, it requires that QED and electroweak corrections be included in all predictions for LHC.

682 Three elements are needed here. Electroweak corrections at NLO order are generally comparable to NNLO
683 QCD corrections, so both types of corrections should be included, together with $\mathcal{O}(\alpha_w \alpha_s)$ terms if possible.
684 Electromagnetic corrections to hadronic reactions cannot be consistently included without a set of PDFs
685 derived from formulae that include NLO QED corrections. This requires a nontrivial modification of PDF
686 fitting programs in order to introduce a photon PDF for the proton. Photon-induced reactions can contribute
687 to LHC processes at the few-percent level, increasing to the 10% level at higher pp energies.

688 Finally, at energies of a TeV and above, electroweak Sudakov effects, negative corrections to two-particle
689 production proportional to $\alpha_w \log^2 s/M_W^2$, can become important. These are 10% corrections for Drell-
690 Yan processes producing 3 TeV dilepton systems. At higher energy pp colliders, these double logarithmic
691 corrections must be resummed systematically.

692 1.6.4 High-precision calculation

693 In the past decade, a revolution in calculational technique has made it possible to derive formulae at NLO
694 for the QCD cross sections for complex multiparton processes such as $pp \rightarrow W + 4$ jets and $pp \rightarrow t\bar{t} + 2$ jets.
695 This has reduced the size of the theoretical errors in these cross sections from order 1 to 10-20%. Methods
696 are now being developed to evaluate general 2-parton processes and even some 3-parton production processes
697 to NNLO, to reduce these theoretical errors to the few-percent level.

698 We have already made reference to NNLO calculations of $t\bar{t}$ and 2-jet production. A very important target
699 here is the cross section for Higgs boson production in association with one or more jets. Many Higgs
700 measurements at the LHC include jet vetoes to control background from $t\bar{t}$ production and other sources,
701 so explicit accounting for emitted jets is necessary. These cross sections often require terms to NNLO for
702 stable summation of the perturbation series.

703 Beyond the fixed-order perturbation theory, many other aspects of higher-order computation remain to be
704 understood. NNLO computations often display large logarithms, which should be systematically resummed.
705 The merging of Monte Carlo programs with NLO QCD calculations is incompletely understood, and new
706 difficulties arise at NNLO. We are optimistic that Higgs boson production and other QCD processes can be
707 computed to few-percent accuracy, but many challenges remain.

708 1.6.5 The Message

709 The conclusions of the QCD working group can be summarized as follows:

- 710 1. Improvements in PDF uncertainties are required. There are strategies at LHC for these improvements.
 711 QED and electroweak corrections must be included in PDFs and in perturbative calculations.
- 712 2. An uncertainty in α_s of order 0.1% may be achievable through improvements in lattice gauge theory
 713 and precision experiments.
- 714 3. Advances in all collider experiments, especially on the Higgs boson, require continued advances in
 715 perturbative QCD.

716 Experiments on QCD give information on the Particle Physics Questions # 1, 2, 8, 9 listed in the Snowmass
 717 Summary [31].

718

719 1.7 Fully Understanding the Top Quark

720

721 The top quark is the heaviest quark and, indeed, the heaviest elementary particle known today. Its large
 722 mass give it the strongest coupling to the Higgs boson and to other possible particles of the Higgs sector.
 723 The mass of the top quark seems to be anomalously large—though it is sometime argued that it is the masses
 724 of all other quarks and leptons that are anomalously small. For all of these reasons, the top quark merits
 725 thorough experimental investigation.

726 The Tevatron experiments that discovered the top quark produced, in all, about 100,000 of these particles.
 727 The LHC experiments have already produced 10 million and aim for many billions of top quarks by the end
 728 of the HL-LHC. Future lepton colliders will bring new precision tools to the study of the top quark. In this
 729 section, we will discuss what can be learned from these observations. More details on all of these topics can
 730 be found in the working group report [39].

731 1.7.1 Top quark mass

732 Like α_s discussed in the previous section, the top quark mass is a crucial input parameter for many SM
 733 predictions. It is already the most accurately known quark mass; a 2 GeV uncertainty on this quantity
 734 corresponds to a measurement with 1% precision. An accurate top quark mass is needed for precision
 735 electroweak fits, with an error of 600 MeV on the top quark mass yielding, for example, an error of 5 MeV
 736 in M_W . The top quark mass is also an important input to the question of ultimate vacuum stability in the
 737 SM [40]. Figure 1-7 shows the suspiciously marginal position of the measured Higgs boson mass and that
 738 of the top quark. Clearly more precision on the latter would help to elucidate whether this is another hint
 739 of strange behavior in the Higgs sector.

740 The top quark mass is most precisely defined as an \overline{MS} quantity, evaluated most conveniently at the \overline{MS} top
 741 quark mass value itself. However, experimental determinations of the top quark mass are typically done by
 742 kinematic fitting to templates, with poorly controlled errors from Monte Carlo modeling and hadronization.
 743 Thus, the precision determination of the top quark mass contains a double challenge, first, to give a definition
 744 of the top quark mass that can be cleanly related to the \overline{MS} mass, and then to measure that quantity
 745 accurately.




Figure 1-7. Regions of metastability and instability of the Higgs potential in the SM, as the top quark and Higgs boson masses are varied, from [40].

746 One solution to this challenge at the LHC is an idea from CMS[41] to measure the top quark mass from the
 747 endpoint of the distribution of the mass $m(b\ell)$ in top quark pair production, where ℓ is an isolated lepton and
 748 b is a b -tagged jet defined by anti- k_T or a similar prescription. The distribution of $m(b\ell)$ can be computed in
 749 QCD perturbation theory in terms of the perturbative pole mass, which can be related to the \overline{MS} mass with
 750 an error of the order of 200 MeV. The endpoint feature in the distribution is sharp and strongly dependent
 751 on m_t . We expect that this method can reach a total uncertainty of 500 MeV with the statistics of the
 752 HL-LHC. Other methods for measuring the top quark mass at the LHC are discussed in [39].

753 At lepton colliders, the cross section for $t\bar{t}$ production near the threshold has a distinctive rise sensitive to the
 754 position of the lowest (unstable) $t\bar{t}$ bound state. Extensive theoretical work has evaluated this cross section
 755 to NNLO, with resummation of all large logarithms. The threshold position can be measured to 35 MeV at
 756 the ILC and somewhat better at TLEP and muon colliders. The conversion to the \overline{MS} mass gives a total
 757 uncertainty of about 100 MeV. This very accurate value of m_t is well matched to the precision electroweak
 758 programs at lepton colliders described in Section 1.5.

759 1.7.2 Strong and electroweak couplings

760 The production and dynamics of top quark pairs at colliders offers many opportunities to test the strong and
 761 electroweak couplings of these particles. At hadron colliders, the dominant pair production mechanism is
 762 through QCD. The current agreement between the predicted and measured values verifies that the absolute
 763 strength of the QCD coupling to the top quark is equal to the value of α_s measured elsewhere to about 3%
 764 accuracy.

765 Changes in the form of the top quark coupling to gluons might be induced by new resonances associated with
 766 top quark compositeness. Possible magnetic or electric dipole couplings can be probed from the kinematics
 767 of top final states to better than 1% at the LHC with 300 fb^{-1} . Though it is difficult to measure the absolute
 768 size of the top quark width at a hadron collider, the W helicity fractions in top quark decay are sensitive
 769 to modifications of the top quark coupling to W , with similar sensitivity. The cross section for single top
 770 production provides a measure of the CKM matrix element V_{tb} which should reach an accuracy of 2.5% at
 771 300 fb^{-1} . Couplings of the top quark to the photon and Z are constrained by measurements of radiation
 772 from a $t\bar{t}$ state. The HL-LHC is expected to reach sensitivities of a few percent for the photon couplings and
 773 15-20% for the Z couplings.

774 At lepton colliders, $t\bar{t}$ pairs are produced through virtual photons and Z s, with large interference effects that
 775 depend on the beam polarization. The ILC and CLIC, which can take advantage of large beam polarization,
 776 expect to reach sensitivities below the 1% level for both photon and Z couplings. Randall-Sundrum models
 777 and other models with Higgs compositeness predict shifts of the Z couplings to $t\bar{t}$ at the few percent level;
 778 these effects could potentially be discovered in the linear collider programs.

779 1.7.3 Rare decays

780 The large samples of top quarks available at the LHC allow deep searches for flavor-changing top quark
 781 decays. Neutral current decays of the top quark such as $t \rightarrow \gamma c$ or $t \rightarrow gc$ are utterly negligible in the SM,
 782 with predict branching ratios smaller than 10^{-12} . These decays can appear with branching ratios as large as
 783 10^{-4} in models with an extended Higgs sector or R-parity violating supersymmetric couplings that bring in
 784 two structures of flavor mixing. Searches for these decays at the HL-LHC can reach branching ratio limits
 785 below 10^{-5} .

786 Lepton colliders can also access these flavor changing couplings in single top production, for example, through
 787 $\gamma^*, Z^* \rightarrow t\bar{c}, t\bar{u}$. Searches for these processes can reach sensitivities close to 10^{-4} even in experiments at
 788 250 GeV, below the $t\bar{t}$ threshold, and below 10^{-5} in the full ILC program at 500 GeV.

789 More details on the specific estimates for each possible neutral current coupling can be found in [39].

790 1.7.4 Searches for new particles related to the top quark

791 The motivation that we have given for new particles at the TeV scale in Section 1.2.2 directly implies the
 792 presence of exotic partners of the top quark. Examples of these particles are scalar top quarks in models of
 793 SUSY and Kaluza Klein excitations of top quarks in models with extra space dimensions. Searches for these
 794 particles have been a very high priority in the LHC program and will continue to be pursued intensively as
 795 more data accumulates.

796 Searches are designed individually for each type of exotic particle. The most powerful searches make use
 797 of the fact that top quarks resulting from the decay of the particle have different polarizations than those
 798 typically produced in SM pair-production. This is reflected in the kinematic distributions of the $t\bar{t}$ final
 799 states. For particles with masses of 1 TeV and above, the preferred method for identification of final-state
 800 top quarks is as single jets with high jet mass and a 3-jet substructure [42]. This “boosted top” identification
 801 is quite insensitive to the pileup associated with high luminosity.

802 Top squarks in SUSY might be tied to the general mass scale of supersymmetric particles, but the naturalness
 803 arguments we have given in Section 1.2.2 indicate that they might be the lightest colored supersymmetric
 804 partners. The LHC experiments have searched extensively for direct pair-production of top squarks that
 805 decay to the lightest supersymmetric particle $\tilde{\chi}^0$ through $\tilde{t} \rightarrow t\tilde{\chi}^0$ and $\tilde{t} \rightarrow b\tilde{\chi}^+$. Current searches exclude
 806 a top squark up to about 650 GeV in the limit of light electroweak superpartners. The sensitivity should
 807 advance to about 1 TeV at 14 TeV and 300 fb $^{-1}$, and to 1.2 TeV with 3000 fb $^{-1}$.

808 In composite Higgs and extra dimensional models, the expected partners of the top quark are fermions
 809 with vectorlike couplings. The searches for these particles are similar to those for fourth-generation quarks,
 810 but they involve more complex decay patterns, with $T \rightarrow Wb$, $T \rightarrow tZ$ and $T \rightarrow th$. Searches for these
 811 particles that are comprehensive with respect to the decay mode currently exclude vectorlike top partners

812 up to masses of about 650 GeV. The 14 TeV stages of the LHC will be able to discover these particles at
813 masses of sensitivity to 1.3 TeV for 300 fb⁻¹ and to 1.6 TeV for 3000 fb⁻¹.

814 Composite Higgs models also typically include resonances in the multi-TeV mass region that decay prefer-
815 entially to $t\bar{t}$. Randall-Sundrum models, for example, predict a resonance at a mass of a few TeV decaying
816 with high top quark polarization to $t_R\bar{t}_L$. The boosted top identification described above was developed
817 for the problem of discovering such states and is indeed expected to be very effective. Applying the same
818 methods to larger data sets, we expect a sensitivity to such resonances up to 4.5 TeV for the 14 TeV LHC
819 with 300 fb⁻¹ and up to 6.5 TeV with 3000 fb⁻¹.

820 Additional examples of new particle searches involving top quarks are described in [39].

821 1.7.5 The Message

822 The conclusions of the Top Quark working group can be summarized as follows:

- 823 1. The top quark is intimately tied to the problems of symmetry breaking and flavor.
- 824 2. Precise and theoretically well-understood measurements of top quark masses are possible both at LHC
825 and at e^+e^- colliders, in each case, matching the needs of the precision electroweak program.
- 826 3. New top couplings and new particles decaying to top play a key role in models of Higgs symmetry
827 breaking. LHC will search for the new particles directly. Linear collider experiments will be sensitive
828 to predicted deviations from the SM in the top quark couplings.

829 Experiments on the top quark give information on the Particle Physics Questions # 1, 2, 4, 8, 9 listed in
830 the Snowmass Summary [31].

831

832 1.8 The Path Beyond the Standard Model - New Particles, Forces, 833 and Dimensions

834

835 Models of new physics associated with the TeV mass scale contain a wide variety of new particles. These
836 include the particles of an extended Higgs sector discussed in Section 1.4.5 and partners of the top quark
837 discussed in Section 1.7.4. More generally, many of the schemes discussed in Section 1.2.2 for explaining Higgs
838 condensation are based on far-reaching principles that require a spectroscopy of new particles containing
839 heavy partners for all SM particles. This includes additional strongly interacting particles, particles with
840 only electroweak interactions, and new vector bosons. Some of these particles may have lifetimes long
841 enough that their decays are not prompt in a collider experiment. One or more of these particles could be
842 the constituents of the cosmic dark matter.

843 New particles could also introduce new flavor-changing interactions. Observation of such interactions of
844 new particles can complement searches for flavor and CP violation in rare processes. This subject will be
845 discussed in Section 1.9.

846 The LHC experiments have been able to search for new particles very robustly over a broad range of
 847 properties. In this section, we will discuss how higher energies and luminosities at hadron colliders and new
 848 capabilities of lepton colliders will extend these searches.

849 Models of Higgs condensation and other TeV-scale phenomena based on different underlying principles make
 850 qualitatively different predictions for the quantum numbers and mass relations of the new particle spectrum.
 851 Thus, the first discovery of a new particle beyond the Standard Model will define a direction for an extensive
 852 research program, one that will be carried out over decades with multiple, complementary, experiments.
 853 In this section, we will emphasize the comparative reach of proposed collider programs to make this first
 854 discovery. Examples of the consequences of such a discovery will be given in Section 1.11. We will have room
 855 to discuss only a limited number of examples. The full range of searches for new particles accessible to TeV
 856 energy experiments is described in the New Particles and Forces working group report [43].

857 The dependence of search reach on luminosity deserves comment. Away from kinematic limits for a given
 858 collider energy, parton-parton luminosity functions scale such that increasing the parton-parton center of
 859 mass energy by a factor 2 decreases the luminosity by a factor 10. This rule, which implies that a factor of 10
 860 in luminosity increases search reach by a factor of 2 in mass, must break down at masses near the kinematic
 861 limit. At the 14 TeV LHC, the reach increase falls off from the canonical factor of 2 for pair-produced
 862 particles with masses well above 1 TeV.

863 1.8.1 New Vector Bosons

864 We begin by discussing particles that show up in collider experiments as distinct resonances. An example is a
 865 color-singlet vector boson associated with an extension of the Yang-Mills symmetry group beyond that of the
 866 SM. Such bosons are required in many contexts, including models with left-right symmetric weak interactions
 867 at high energy, models of the Higgsino mass in SUSY, and models with extra dimensions. Models of Higgs
 868 composite structure often require breaking of a larger gauge group to the SM symmetry group.

869 Searches for these bosons are conducted at hadron colliders by looking for narrow dilepton resonances. A
 870 typical benchmark is sensitivity to the “sequential SM” Z' , a boson with the couplings of the Z but with a
 871 higher mass. Current results from the LHC require the mass of such a particle to be above 2.5 TeV. With 14
 872 TeV, it will be possible to discover such a resonance at 4.5 TeV, for 300 fb^{-1} , and at 7 TeV, for 3000 fb^{-1} .
 873 The value of the production cross section and the leptonic forward-backward asymmetry (with respect to
 874 the direction of production) give information on the couplings of the Z' . At higher pp energies, the discovery
 875 reach increases to 12 TeV for 33 TeV and 30 TeV for the 100 TeV VLHC.

876 Lepton colliders are sensitive to new vector bosons that interfere with the s -channel virtual photon and Z in
 877 two-fermion production $\ell^+\ell^- \rightarrow f\bar{f}$. The reach for discovery of a sequential Z' at the ILC at 500 GeV is also
 878 about 7 TeV and scales proportional to the center of mass energy for higher energy colliders. Measurements
 879 of the Z' signal with two possible beam polarizations and with individual lepton and quark final states gives
 880 a large amount of information toward the identification of the quantum numbers of the boson.

881 1.8.2 Supersymmetry

882 Searches for supersymmetry (SUSY) encompass a wide range of strategies aimed at different particles of the
 883 SUSY spectrum.




Figure 1-8. Projections for coverage of the pMSSM 19-parameter model space in searches at the LHC at 300 fb^{-1} (left) and 3000 fb^{-1} (right), shown in the plane of gluino mass versus lightest particle mass. Black is complete exclusion; green is exclusion of 50% of the pMSSM model points. From [47].

884 The most generic searches assume that supersymmetric partners of the SM particle carry a conserved
 885 quantum number, called R-parity. If the lightest supersymmetric particle is neutral, it will typically be
 886 weakly interacting and will not be observed in a collider detector. Events are then characterized as containing
 887 several hadronic jets, associated with decay to the lightest particle plus missing transverse momentum. The
 888 results of these analyses are parametrized by limits on the gluino mass and on a squark mass, assumed
 889 common to all squark flavors. Current LHC results exclude such events up to gluino masses of 1.0 TeV and,
 890 independently, squark masses of 1.3 TeV. For the future stages of the LHC, we expect to be able to discover
 891 such events up to gluino masses of 1.9 TeV and squark masses of 2.3 TeV with 300 fb^{-1} , and to 2.3 TeV
 892 and 2.7 TeV with 3000 fb^{-1} . This reflects more than a factor 2 in increased search power at the 300 fb^{-1}
 893 stage, and another 20% with the additional factor 10 in luminosity. The gluino discovery reach increases to
 894 4.8 TeV for a 33 TeV pp collider and to 10.2 TeV for the VLHC.

895 It is possible that the first signal of SUSY would not be given by the generic search just described, but would
 896 require a more specialized analysis. Special search techniques are needed in models in which mass gaps in the
 897 SUSY spectrum are relatively small (“compressed spectrum”), so that hard jets are not emitted in particle
 898 decays, and models in which only the partners of top quarks, or perhaps, only color-singlet supersymmetric
 899 particles are produced at accessible energies. Such models can be highly motivated. A compressed spectrum
 900 is needed in many models of supersymmetric dark matter particles to allow “coannihilation” to produce
 901 the correct dark matter density [44]. They are also needed in models in which only the specific particles
 902 restricted by the naturalness bounds in Section 1.2.2 lie below 2 TeV. In such models, the first signal of
 903 SUSY would come from direct top squark pair production or gluino pair production with decay to heavy
 904 flavor. Reach estimates for top squark pair production were given above in Section 1.7.4.

905 Models in which SUSY discovery is more difficult at the 8 TeV LHC thus benefit more from the increase in
 906 luminosity to the HL-LHC. Models for which the first signal of SUSY would be the partners of W and Z
 907 bosons can be searched for at the 14 TeV LHC, with discovery expected up to masses of about 500 GeV.
 908 The factor of 10 luminosity increase to HL-LHC increases the reach by a factor of 2 in the analyses [45, 46],
 909 consistent with the argument given above.

910 Another way to look at this issue is shown in Fig. 1-8. The figures shows a survey of a large number of SUSY
 911 models [47] plotted in the plane of gluino mass versus lightest superparticle mass. The color-coding gives
 912 the fraction of models excluded by LHC searches, at 300 fb^{-1} on the left, and at 3000 fb^{-1} on the right.

913 The general shift of the boundary by about 30% is accompanied by a removal of exceptions to the right of
914 this boundary.

915 One more exception should be noted. The supersymmetric partners of the Higgsino automatically have small
916 mass splitting of a few GeV. The direct pair-production of these particles through electroweak interactions
917 is essentially invisible at the LHC, except through searches for initial-state radiation plus invisible particles,
918 described in Section 1.8.4. In a scenario in which these are the lightest supersymmetric particles, the ability
919 of lepton colliders to be sensitive to very small energy depositions in decay would be crucial to observe and
920 study these particles. Studies of Higgsino pair production at the ILC as described in [48].

921 1.8.3 Long lived particles

922 The searches we have described so far assume that all new particles decay promptly. However, there are
923 many models that give exceptions to this. ATLAS and CMS have carried out dedicated searches for tracks
924 associated with long-lived massive particles and for particles decaying in the detector, perhaps out of time
925 with the bunch crossings. Current limits are stronger than those in searches for promptly decaying particles.
926 For example, ATLAS places limits of 310 GeV on a tau slepton, 600 GeV on a top squark, and 985 GeV
927 on a gluino. Should such long-lived particles exist, the LHC detectors trap a sample of them for detailed
928 studies of their decay modes and lifetimes.

929 1.8.4 Dark matter

930 The signature of jets plus missing transverse momentum discussed above for SUSY applies to a broader class
931 of theories. In Section 1.2.3, we introduced WIMP dark matter as a general motivation for new particles
932 with TeV scale masses. Any model with a new TeV spectroscopy characterized by a new quantum number
933 can give rise to a dark matter candidate particle. The requirements are that the lightest new particle is
934 neutral and that the quantum number is conserved sufficiently that this particle is stable over the age of
935 the universe. Heavier states carrying QCD color will decay to this lightest particle, producing events with
936 jets and missing transverse momentum. If the partners of quarks are fermionic rather than bosonic, the
937 production cross section will be higher. The generic SUSY search described above can then be interpreted
938 as a robust search for models predicting a TeV spectroscopy and dark matter.

939 It is possible that the heavier states of the TeV spectrum are out of reach kinematically. Then the discovery of
940 dark matter would require the observation of direct pair-production of the essentially invisible dark matter
941 particles. This can be done using the fact that the production of particles in a hard scattering process
942 sometimes also produces gluons or photons radiated from the initial-state particles. If the particles mediating
943 the pair-production reaction are heavy enough, this initial-state radiation can be at higher momentum than,
944 and, thus, distinguishable from, the ordinary particle production in typical collisions.

945 Reach estimates for the discovery of dark matter pair-production have been studied systematically in [49],
946 using an effective operator formalism to describe the coupling of dark matter to SM particles. This formalism
947 also allows cross sections measured at colliders to be related to rates for direct and indirect detection of dark
948 matter. The paper [49] also included limits on an explicit model with a lighter Z' mediator. Some results
949 of the effective field theory analysis, comparing limits on the dark matter-nucleon cross section from LHC
950 and higher energy pp colliders to limits from direct detection, are shown in Fig. 1-9. It is noteworthy that




Figure 1-9. Comparison of 95% confidence limits on the WIMP pair annihilation cross section in models of dark matter, analyzed in an effective operator formalism, for current and proposed facilities. The collider limits are constraints on the operator coefficient from searches for missing energy events; The limits from gamma ray telescopes are constraints from searches for dark matter annihilation in galaxies of the local group. From [49].

951 the VLHC can place limits on the dark matter particle mass above 1 TeV, close to the unitarity limit for
 952 thermal production of such a particle in the early universe.

953 1.8.5 The Message

954 The conclusions of the New Particles and Forces working group can be summarized as follows:

- 955 1. TeV mass particles are needed in essentially all models of new physics. The search for them is
 956 imperative.
- 957 2. LHC and future colliders will give us impressive capabilities for this study. Future programs target
 958 new physics at the all-important TeV scale, as can be seen in Fig. 1-10.
- 959 3. The search for TeV mass particles is integrally connected to searches for dark matter.

960 Experiments on new particles and forces give information on the Particle Physics Questions # 1, 2, 3, 4, 5,
 961 8, 9 listed in the Snowmass Summary [31].

962

963 1.9 Flavor Mixing and CP Violation at High Energy

964




Figure 1-10. Examples of 95% confidence upper limits for new particle searches at proposed pp and e^+e^- colliders.

965 The explanation of Higgs condensation may or may not give insight into the theory of flavor. It is a mystery
 966 why quarks and leptons have a hierarchical mass spectrum, why the weak interactions are not diagonal in
 967 flavor and violate CP. In the SM, these features are parametrized by the fermion-Higgs Yukawa couplings.
 968 The idea that flavors are distinguished only through terms of the structure of the Yukawa couplings is called
 969 “Minimal Flavor Violation.”

970 Models of new physics at the TeV scale introduce a large number of new couplings. These in principle
 971 can be proportional to flavor-violating couplings with completely new structures. Such structures are not
 972 needed to build a model of Higgs condensation. In fact, many specific models—most notably, SUSY with
 973 gauge-mediated breaking [56]—are flavor-blind. It is possible also that some principle, analogous to the GIM
 974 mechanism, requires that new flavor couplings are related to the Yukawa couplings. In these cases, there are
 975 no new flavor-changing effects beyond the SM arising from the TeV scale.

976 However, it is also possible that flavor couplings among new particles have a different pattern. Such couplings
 977 can generate rare flavor-changing weak decays. They can also affect the phenomenology of the new particles
 978 themselves, requiring new strategies for searches. In this section, we discuss examples of models of this type.
 979 There are many possibilities, only a few of which can be discussed here. A more complete catalog is given
 980 in the working group report [50].

1.9.1 SUSY with Flavor-Dependent Soft Masses

In Section 1.8, we discussed reach estimates for models of TeV spectroscopy that were either blind to flavor or singled out only the third generation. More general forms of the new particle spectrum are allowed. The most important difficulties with flavor observables come when squarks with the same gauge quantum numbers, *e.g.*, the partners of d_R and s_R , have different masses. But there is little difficulty in giving the partners of d_R and d_L smaller masses than those of the other squarks. This weakens the experimental limits on the lightest squark mass. In fact, it is possible, with other mechanisms to suppress flavor effects of new physics, to allow only the partner of c_R to be light. This has been explored in detail for SUSY models [51, 52]. The current limit on the charm squark in such models is about 300 GeV.

1.9.2 R-parity Violating SUSY

It is possible that the R-parity conservation law that should keep the lightest SUSY partner stable is violated by new interactions. These interactions necessarily have a complex structure in flavor. One possible form of the R-parity violating interaction is

$$\lambda_{ijk}^1 U_i D_j D_k , \quad (1.17)$$

where U_i , D_i are the right-handed quarks or their squark partners and $i = 1, 2, 3$ is the generation number. This violates baryon number, allowing a squark to decay to two antiquarks. The interaction must be antisymmetric in color, and this requires it also to be antisymmetric in the flavors of the two down-type quarks. Other possible R-parity violating interactions have the forms

$$\lambda_{ijk}^2 L_i L_j \bar{E}_k , \quad \lambda_{ia}^3 L_i H_a \quad (1.18)$$

where L_i , \bar{E}_i are left-handed leptons or antileptons or their slepton partners and H_a is a Higgs or Higgsino field. These interactions violate lepton number. The coefficients of these operators are usually taken to be small to avoid unwanted flavor-changing rare decays. In particular, either the operator (1.17) or the lepton number violating operators must be highly suppressed to avoid rapid proton decay.

In R-parity violating models with the operator (1.17) only, SUSY decay chains typically end with the lightest SUSY particle decaying to jets. These jets must have a nontrivial flavor structure, possibly with a b jet always included.

R-parity violation with the Higgs operator in (1.18) can produce neutrino masses through the ‘‘Type III seesaw’’: A neutrino converts to a Higgsino, which then converts back to an antineutrino, possibly of a different flavor. The difference between the quark and lepton flavor mixing patterns is explained by the statement that the mass matrices come from unrelated operators. In such models, the branching ratios of the Higgsino are related to the neutrino mixing angles, and this relation can be confirmed by direct measurement [53]. In more general seesaw models of neutrino mass, it is possible that the seesaw scale could be the TeV scale, setting up other relations between TeV mass neutral leptons and the neutrino mixing matrix [54].

1.9.3 Models with Electroweak Baryogenesis

To explain the asymmetry between the numbers of baryons and antibaryons in the universe, a new source of CP violation is needed beyond that of the CKM phase. One natural place to look for this is in an

1016 extended Higgs sector. A second requirement is that the electroweak phase transition be first-order. Then
 1017 the transition takes place through the growth of bubbles, with top quarks scattering from the bubbles
 1018 because they are massless outside and massive inside. A CP phase in the Higgs sector makes this scattering
 1019 asymmetric between matter and antimatter. Both criteria can be satisfied in models with multiple Higgs
 1020 fields [55]. Measurements of the Higgs self-coupling and of CP violation in Higgs decays, described in
 1021 Sections 1.4.2 and 1.4.3 above, test models of this type.

1022 1.9.4 The Message

1023 The conclusions of the Flavor and CP working group can be summarized as follows:

- 1024 1. TeV mass particles may or may not introduce couplings with new types of flavor violation. These
 1025 possible new couplings affect the search methods for new particles, in many cases, requiring new
 1026 strategies.
- 1027 2. The search for new particles is integrally connected to searches for rare flavor changing decays.

1028 Experiments on flavor associated with new particles give information on the Particle Physics Questions #
 1029 1, 2, 3, 4, 5, 8, 9 listed in the Snowmass Summary [31].

1030 1.10 Scientific Cases for Future Colliders

1031 In the previous sections, we presented the physics opportunities for the next steps in the Energy Frontier in
 1032 terms of individual particles and research areas under study. It is also interesting to assemble these topics
 1033 in terms of the experimental program that each accelerator in the list given in Section 1.3.2 will provide.
 1034 Integrating over the topics in this way, we see that many of these proposed accelerators have very substantial
 1035 physics programs that will explore the TeV energy scale across a broad range of measurements.

1036 In this section, we present the cases for the various accelerators as if each accelerator stood on its own,
 1037 with no further physics discoveries between now and the time that it begins operation. However, one should
 1038 always keep in mind the possibility of discoveries would open up the study of physics beyond the SM. We
 1039 have argued already that the likelihood of the discovery of new particle is very high even for the coming runs
 1040 of the LHC at 14 TeV up to 300 fb^{-1} . Such a discovery would need to be followed up by further exploration
 1041 that would benefit from accelerators with complementary capabilities or higher energy. This might, in the
 1042 end, be the most important benefit of building the accelerators that come later in the timeline below. We
 1043 will expand on this idea in Section 1.11.

1044 1.10.1 LHC in This Decade: 300 fb^{-1}

1045 First of all, we emphasize the many opportunities that will be provided by the coming run of the LHC
 1046 at 14 TeV. Operation of the LHC at 14 TeV up through the 300 fb^{-1} with the Phase 1 upgrades offers a
 1047 tremendous increase in the power of new particle searches, close to a factor 2 in mass in most channels.
 1048 Many of these searches, for example, the search for the gluino almost to 2 TeV and the search for vectorlike
 1049 top partners above 1 TeV, access ranges of the masses that are strongly motivated in models of Higgs
 1050 condensation.

1051 This impressive capability is only one aspect of a broad program that will be carried out at the LHC over
1052 the next ten years. Its features include:

- 1053 1. Clarification of Higgs boson couplings, mass, spin, CP to the 10% level.
- 1054 2. First direct measurement of top-Higgs boson couplings
- 1055 3. Precision W boson mass measurement below 10 MeV.
- 1056 4. First measurements of VV scattering.
- 1057 5. Theoretically and experimentally precise top quark mass to 600 MeV.
- 1058 6. Measurement of top quark couplings to gluons, Z and W bosons, and photons with a precision
1059 potentially sensitive to new physics—a factor 2–5 better than today.
- 1060 7. Search for top squarks and top partners and $t\bar{t}$ resonances predicted in models of composite top, Higgs.
- 1061 8. New generation of PDFs with improved gluon and antiquark distributions.
- 1062 9. Precision study of electroweak cross sections in pp collisions, including a new photon PDF.
- 1063 10. Extension by a factor 2 in the sensitivity to new particles: SUSY, Z , top partners—key ingredients for
1064 models of the Higgs potential—and the widest range of possible TeV-mass particles.
- 1065 11. Deep ISR-based searches for dark matter particles.

1066 1.10.2 High-Luminosity LHC: 3 ab^{-1}

1067 The second high luminosity running of the LHC is referred to as the “Phase 2” upgrade period with
1068 instantaneous luminosities of $5 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$. This running with $3,000 \text{ fb}^{-1}$ of accumulated data truly
1069 inaugurates the high-precision electroweak era at LHC with few percent precision for most Higgs boson
1070 couplings as well as the 5 MeV threshold in M_W mass determination.

- 1071 1. The precision era in Higgs boson couplings begins, with sensitivities to 2-10%, and 1% for the ratio of
1072 $\gamma\gamma$ and ZZ couplings.
- 1073 2. Measurement of rare Higgs boson decays, $\mu^+\mu^-$ and $Z\gamma$, with 100 M Higgs bosons.
- 1074 3. First evidence of the Higgs boson self-coupling.
- 1075 4. Powerful searches for extended Higgs bosons.
- 1076 5. Precision W mass to $\pm 5 \text{ MeV}$.
- 1077 6. Precise measurements of VV scattering with access to Higgs sector resonances.
- 1078 7. Precision top mass to $\pm 500 \text{ MeV}$.
- 1079 8. Deep study of rare, flavor-changing, top couplings with 10G tops.
- 1080 9. Search for top squarks and partners in models of composite top quarks and Higgs bosons in the expected
1081 range of masses.
- 1082 10. Further improvement of q , g , and γ PDFs to higher x and Q^2 .

- 1083 11. A 20-40% increase in mass reach for generic new particle searches which can be as much as a 1 TeV
1084 step in mass reach.
- 1085 12. Extension by a factor of 2 in the mass reach for particles produced by the electroweak interactions.
- 1086 13. Any discovery at LHC — or in dark matter or flavor searches — can be followed up.

1087 1.10.3 ILC, up to 500 GeV

1088 The ILC would run at 250 GeV, 350 GeV, and 500 GeV, in a program that could begin as early as the
1089 second half of the next decade. It will study the properties of the Higgs boson, the top quark, and possibly
1090 also newly discovered particles, in very fine detail.

- 1091 1. Tagged Higgs boson study in $e^+e^- \rightarrow Zh$: model-independent Higgs boson width and branching ratio
1092 measurements, direct study of all Higgs decay modes, including invisible and exotic decays.
- 1093 2. Model-independent Higgs boson couplings with percent-level precision necessary to probe for new
1094 physics beyond the reach of LHC direct searches.
- 1095 3. Higgs boson CP studies in fermionic channels (e.g., $\tau^+\tau^-$).
- 1096 4. Giga-Z program for EW precision, W boson mass to 4 MeV and beyond.
- 1097 5. Improvement of triple vector boson couplings by a factor 10, to an accuracy below expectations for
1098 models with Higgs sector resonances.
- 1099 6. Theoretically and experimentally precise top quark mass to ± 100 MeV.
- 1100 7. Sub-% measurement of top couplings to γ and Z , with accuracy well below expectations in models of
1101 composite top quarks and Higgs bosons.
- 1102 8. Search for rare top couplings in $e^+e^- \rightarrow t\bar{c}, t\bar{u}$.
- 1103 9. Improvement of α_s from the Giga-Z program.
- 1104 10. Unambiguous search capability for new particles in LHC blind spots – Higgsino, stealth stop, com-
1105 pressed spectra, WIMP dark matter.

1106 1.10.4 ILC at 1 TeV

1107 In the farther future, the extension of ILC to 1 TeV will access additional Higgs boson reactions for precision
1108 study and, possibly, also reach new particle thresholds.

- 1109 1. Precision measurement of the Higgs boson coupling to top, to 2%.
- 1110 2. Measurement of the Higgs boson self-coupling, to 13%.
- 1111 3. Discovery of any extended Higgs boson states coupling to the Z , up to 500 GeV.
- 1112 4. Improvement in precision of triple gauge boson couplings by a factor 4 over 500 GeV results.
- 1113 5. Model-independent search for new particles with coupling to γ or Z to 500 GeV

- 1114 6. Search for Z using $e^+e^- \rightarrow f\bar{f}$ to masses of about 5 TeV, a reach comparable to LHC for similar
1115 models. Multiple observables for diagnostics of the Z couplings.
- 1116 7. Any discovery of new particles dictates a lepton collider program: search for electroweak partners, 1%
1117 precision mass measurement, the complete decay profile, model-independent measurement of cross
1118 sections, branching ratios and couplings with polarization observables, search for flavor and CP-
1119 violating interactions

1120 1.10.5 CLIC: 350 GeV, 1 TeV, 3 TeV

1121 Extremely high energies in e^+e^- collisions will likely require technologies beyond that envisioned for the
1122 ILC. CLIC is proposed as an e^+e^- collider capable of multi-TeV energies. It would probe Higgs boson
1123 self-couplings and exotic scattering of both standard model particles and any new particles found or hinted
1124 at in earlier machines.

- 1125 1. Precision measurement of the Higgs boson coupling to top, to 2%.
- 1126 2. Measurement of the Higgs boson self-coupling to 10%.
- 1127 3. Discovery of any extended Higgs boson states coupling to the Z , up Discovery of any extended Higgs
1128 boson states coupling to the Z , to 500 GeV. to 1500 GeV.
- 1129 4. Improvement in precision of triple gauge boson couplings by a factor 4 over 500 GeV results.
- 1130 5. Precise measurement of VV scattering, sensitive to Higgs boson sector resonances.
- 1131 6. Model-independent search for new particles with coupling to gamma or Z to 1500 GeV: the expected
1132 range of masses for electroweakinos and WIMPs.
- 1133 7. Search for Z using $e^+e^- \rightarrow f\bar{f}$ accessing masses above 10 TeV.
- 1134 8. Any discovery of new particles dictates a lepton collider program, with the elements listed for the
1135 1 TeV ILC.

1136 1.10.6 Muon Collider: 125 GeV, 350 GeV, 1.5 TeV, 3 TeV

1137 A muon collider holds promise as a technique for reaching very high energies in lepton-lepton collisions and
1138 for s -channel production of the Higgs boson and possible additional Higgs states. Studies of the muon collider
1139 are not yet mature, particularly in designing a detector that can overcome the background from decays of
1140 the muons circulating in the ring. However, promising first results were reported at Snowmass.

- 1141 1. Similar capabilities to e^+e^- colliders described above.
- 1142 2. Ability to produce the Higgs boson, and possible heavy Higgs bosons, as s -channel resonances. This
1143 allows a sub-MeV Higgs boson mass measurement and a direct Higgs boson width measurement.

1.10.7 Photon Collider

Another technique for producing Higgs bosons in the s -channel is to convert an electron collider to a photon collider by backscattering laser light from the electron beams. This allows resonance studies at 80% of the electron center-of-mass energy.

1. Production of Higgs or extended Higgs bosons as s -channel resonances, offering percent-level accuracy in $\gamma\gamma$ coupling.
2. Ability to study CP mixture and violation in the Higgs sector using polarized photon beams.

1.10.8 TLEP, Circular e^+e^-

An e^+e^- collider in a very large tunnel offers the possibility of very large integrated luminosity samples at 250 GeV and below, especially if multiple detectors can be used simultaneously.

1. Possibility of up to 10 times higher luminosity than linear e^+e^- colliders at 250 GeV. Higgs boson couplings measurements might still be statistics-limited at this level.
2. Precision electroweak programs that could improve on ILC by a factor 4 in $\sin^2\theta_w$, a factor 4 in M_W , and a factor 10 in M_Z .
3. Theoretically and experimentally precise top quark mass to ± 100 MeV.
4. Search for rare top couplings in $e^+e^- \rightarrow t\bar{c}, t\bar{u}$.
5. Possible improvement in α_s by a factor 5 over Giga- Z , to 0.1% precision.

1.10.9 VHLC, at 100 TeV

One of the ideas at Snowmass that gained momentum through the week was renewed interest in a Very Large Hadron Collider (VLHC). Reinvigorating R&D in a VLHC was a clear recommendation of the New Particles and Forces Group and the conveners.

1. High rates for double Higgs boson production; measurement of Higgs boson self coupling to 8%.
2. Sensitivity to new Higgs bosons and states associated with extended Higgs sectors at 1 TeV.
3. Dramatically improved sensitivity to vector boson scattering and multiple vector boson production.
4. Increased search reach for new particles associated with naturalness—including SUSY particles, top partners, and resonances—by almost an order of magnitude in mass over LHC. This corresponds to two orders of magnitude in fine-tuning.
5. Sensitivity to WIMP dark matter up to TeV masses, possibly covering the full natural mass range.
6. Any discovery at LHC — or in dark matter or flavor searches — can be followed up by more detailed measurements at VLHC, in addition to searches for related higher-mass particles. Both luminosity and energy are relevant.

1.11 Discovery stories

Another way to integrate over the physics topics presented in Sections 1.4–1.9 is to consider the consequences of a discovery of new physics at the LHC later in this decade. We have emphasized, first, that the presence of new particles at the TeV scale is necessary to build a physics explanation of electroweak symmetry breaking, and, second, that the coming run of the LHC, up to 300 fb^{-1} , will improve the depth of searches for new particles by more than a factor of 2. The conclusion from these statements is that the discovery of new physics at the LHC is likely. This means that we should have a plan for following up this discovery and exploring its implications. This program of course depends on the nature of the particle discovered, so a full analysis would be presented as a large number of case studies. We give two examples here for illustration. Further examples of these “discovery stories” are presented in the working group reports.

1.11.1 Well-Tempered SUSY

The New Particles and Forces working group [43] considered in some detail the consequences of a particular SUSY model that could be discovered at the LHC with 300 fb^{-1} . This particular model had a gluino at 1.9 TeV, squarks ranging in mass from 1.3 TeV to 2.6 TeV, and bino and Higgsino states near 200 GeV. The bino and Higgsino were assumed to mix so that the lightest supersymmetric particle (LSP) would be a dark matter particle with the correct thermally generated cosmic density. This dark matter scenario is called the “well-tempered neutralino” [57].

The LHC at 300 fb^{-1} would observe a robust jets plus missing transverse momentum signal. The signal would be dominated by the decay of the lighter squarks to a quark jet plus the unobserved LSP. The mass difference could be measured from kinematic distributions. Assuming that the LSP was light, this would also give an estimate of the squark mass. With the measured cross section, this would favor SUSY over models with fermionic partners.

The HL-LHC would produce some of the heavier squarks and the gluino. Detailed kinematic measurements would identify at least one more mass scale in the spectrum and give further evidence for the SUSY hypothesis. The direct production of electroweak states would not be observed at the LHC, because these states have a compressed spectrum.

A lepton collider with center of mass energy of 500 GeV would be able to pair-produce the Higgsinos and observe their decays to the LSP. Measurement of the polarized cross sections would give information about the quantum numbers of the electroweak states. It would also give an indirect determination of the mass of the electron-type slepton (750 GeV in this model) to 10 GeV. Using this information, it would be possible to evaluate the LSP annihilation cross section and show that it was consistent with that required for a dark matter particle.

Experiments at higher-energy colliders would be needed to discover the heaviest sleptons (at 3.3 TeV) and squarks (at 2.6 TeV). Eventually, the complete SUSY spectrum would be determined, and the data on the mass spectrum could be used to deduce the pattern of SUSY breaking.

1.11.2 $t\bar{t}$ Resonance

An alternative scenario might be based on a Randall-Sundrum model with top and Higgs compositeness. The first evidence of this model would be the discovery of a resonance in pp collisions that decayed to $t\bar{t}$. Such a resonance at 3 TeV would be discovered at the LHC with 300 fb^{-1} . Study of kinematic distributions of the $t\bar{t}$ final state would reveal that the top quarks were highly polarized, a prediction of this model.

The HL-LHC would discover an electroweak singlet top quark partner, and, possibly also, a doublet of quarks with vectorlike coupling to the electroweak interactions. It is possible that very accurate studies of the $t\bar{t}$ spectrum would also reveal the presence of a color-singlet resonance, somewhat below 3 TeV. Its higher-statistics study of the TeV resonance might reveal at decay to $t\bar{c}$ with branching ratio 10^{-3} .

A lepton collider at 500 GeV would observe a significant 3% enhancement of the right-handed top quark coupling to the Z boson. The Higgs boson coupling to $\gamma\gamma$ might be enhanced at the 2% level by the radiative correction from the top quark partners. This would be discovered by combining the high statistics measurement of $BR(\gamma\gamma)/BR(ZZ^*)$ from the HL-LHC with the precise measurement of the Higgs coupling to Z at a lepton collider.

These measurements would give a tantalizing first glimpse of the structure of the underlying composite Higgs model. Experiments at higher energy colliders capable of producing resonances up to 20 TeV in mass would be needed to explore the full structure of the spectrum of states.

1.12 Conclusions

In this report, we have described the future program of research at high-energy colliders and summarized the efforts of six Snowmass 2013 Energy Frontier working groups. The detailed conclusions that we have reviewed come back repeated to a set of four points that deserve special emphasis.

1. The search for new particles with masses of order 1 TeV is a central issue in particle physics. The mysteries associated with the Higgs field and dark matter give compelling arguments for a new particle spectroscopy at the TeV mass scale. Such particles must exist to provide a physics explanation for the Higgs condensation and symmetry breaking. We need to know their nature, and their implications for the laws of physics at very short distances. We have given many examples in which these particles address other major questions of particle physics, including the questions of flavor, dark matter, and the unification of forces.

2. The central capability of high-energy collider experiments is to produce massive elementary particles directly. In Energy Frontier experiments, we observe the W and Z bosons, the top quark, and the Higgs boson as real particles whose production and decay we can study in detail. The same will be true for any new particles that we can create. This is a unique, direct, and powerful method to learn about the laws of physics.

3. There are three essential aspects in the exploration of the physics of the TeV mass scale.

1. We must study the Higgs boson itself in as much detail as possible, searching for signs of a larger Higgs sector and the effects of new heavy particles.

2. We must search for the imprint of the Higgs boson and its possible partners on the couplings of the W and Z bosons and the top quark.

3. We must search directly for new particles with TeV masses that can address important problems in fundamental physics.

1248 4. This program can be realized at accelerators now envisioned to operate in the coming years.

1249 1. We have emphasized the great opportunity that is being provided by the coming operation of the LHC
1250 at 14 TeV. The next stage of the LHC will double the range of searches for new particles and give
1251 similar leaps in capability for other probes of TeV physics.

1252 2. We have projected quantitatively what will be achieved in running the LHC at high luminosity, to
1253 ultimately acquire 3000 fb^{-1} of data per experiment. For some physics topics, the gain is incremental,
1254 but for others, in particular, the precision study of Higgs couplings and the search for new particles
1255 with only electroweak interactions, we move to a qualitatively new level. Looked at in total, this is a
1256 highly motivated physics program.

1257 3. We have listed many essential contributions to the exploration of the TeV scale that can be provided
1258 by lepton colliders. These include precision studies of the Higgs boson, the W and Z bosons, and the
1259 top quark, capable in all cases of discovering percent-level corrections predicted as the effects of TeV
1260 particles. The construction and operation of the ILC in Japan will realize these goals.

1261 4. We have emphasized that the quest to understand the TeV scale will not be finished with the results
1262 of accelerators of the next generation. It is likely that the discovery of new particles at the next
1263 stage of collider physics will open a definite path for exploration to still higher energies. Our study
1264 called attention, in particular, to the capabilities of a VLHC for further exploration of the TeV mass
1265 scale. The journey to still higher energies begins with renewed effort to bring advanced accelerator
1266 technologies to reality.

1267 We emphasized in our introduction that the discovery of the Higgs boson changes everything. This discovery
1268 points to potentially profound modifications of the laws of physics at energies relatively close to those we
1269 now access at accelerators. The quest for new phenomena and the insights they will provide has just begun.

1270 ACKNOWLEDGEMENTS

1271 The Energy Frontier conveners are profoundly grateful to the 26 dedicated conveners of our working groups:
1272 Kaustubh Agashe, Marina Artuso, John Campbell, Sally Dawson, Robin Erbacher, Cecilia Gerber, Yuri
1273 Gershtein, Andrei Gritsan, Kenichi Hatakeyama, Joey Huston, Ashutosh Kotwal, Heather Logan, Markus
1274 Luty, Kirill Melnikov, Meenakshi Narain, Michele Papucci, Frank Petriello, Soeren Prell, Jianming Qian,
1275 Reinhard Schwienhorst, Chris Tully, Rick Van Kooten, Doreen Wackerroth, Lian-Tao Wang, and Daniel
1276 Whiteson.

1277 We are also grateful to the many scientists who assisted us in various ways through the process. We have
1278 benefited from those who have given us technical advice, from those who maintained our contact with the
1279 major collaborations, and from those who took time to help us with individual tasks.

1280 Our technical team was: Jeff Berryhill, Sergei Chekanov, Tom LeCompte, Sanjay Padhi, Eric Prebys, Tor
1281 Raubenheimer, and Eric Torrence. We also thank Markus Klute and Mark Palmer from the Capabilities
1282 group.

1283 Our advisors from the major experiments were: for ATLAS: Ashutosh Kotwal; for CMS: Jim Olsen; for
1284 LHCb: Sheldon Stone; for ILD: Graham Wilson; for SiD: Andy White; from CLIC: Mark Thomson; from
1285 the Muon Collider: Ron Lipton; and for the VLHC: Dmitri Denisov.

1286 Finally, we thank our fellow Snowmass 2013 conveners who asked “tough questions” and answered them.

References

- [1] See, for example, S. Schael *et al.* [ALEPH and DELPHI and L3 and OPAL and SLD and LEP Electroweak Working Group and SLD Electroweak Group and SLD Heavy Flavour Group Collaborations], Phys. Rept. **427**, 257 (2006) [hep-ex/0509008].
- [2] Examples of these mechanics are found, for supersymmetry, in L. E. Ibanez and G. G. Ross, Phys. Lett. B **110**, 215 (1982), L. Alvarez-Gaume, J. Polchinski and M. B. Wise, Nucl. Phys. B **221**, 495 (1983); for Little Higgs modes, in N. Arkani-Hamed, A. G. Cohen, E. Katz and A. E. Nelson, JHEP **0207**, 034 (2002) [hep-ph/0206021]; for extra-dimensional models Y. Hosotani, S. Noda and K. Takenaga, Phys. Lett. B **607**, 276 (2005) [hep-ph/0410193].
- [3] Examples are found, for supersymmetry, in S. Weinberg, Phys. Rev. Lett. **50**, 387 (1983), H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983) [Erratum-ibid. **103**, 099905 (2009)]; for extra-dimensional models, in H. -C. Cheng, J. L. Feng and K. T. Matchev, Phys. Rev. Lett. **89**, 211301 (2002) [hep-ph/0207125].
- [4] Examples are found, for supersymmetry, in S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. D **24**, 1681 (1981); for extra-dimensional models, in K. Agashe, R. Contino and R. Sundrum, Phys. Rev. Lett. **95**, 171804 (2005) [hep-ph/0502222].
- [5] The following discussion oversimplifies an extensive and sophisticated literature on naturalness in supersymmetry. See, *e.g.*, R. Barbieri and G. F. Giudice, Nucl. Phys. B **306**, 63 (1988), B. de Carlos and J. A. Casas, Phys. Lett. B **309**, 320 (1993) [hep-ph/9303291]; J. L. Feng, K. T. Matchev and T. Moroi, Phys. Rev. Lett. **84**, 2322 (2000) [hep-ph/9908309]; M. Papucci, J. T. Ruderman and A. Weiler, JHEP **1209**, 035 (2012) [arXiv:1110.6926 [hep-ph]]; H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev and X. Tata, arXiv:1306.2926 [hep-ph] (Snowmass paper).
- [6] J. Feng, *et al.*, Cosmic Frontier report, Snowmass paper.
- [7] W. Barletta, *et al.*, Snowmass paper.
- [8] ref. on the LHC future
- [9] ref. on High-E LHC at 33 TeV
- [10] ref. on VLHC 100 TeV at CERN
- [11] ref. on VLHC 100 TeV
- [12] J. Brau, P. Grannis, M. Harrison, M. Peskin, M. Ross and H. Weerts, arXiv:1304.2586 [physics.acc-ph], Snowmass paper.
- [13] T. Behnke, *et al.*, arXiv:1306.6327 [physics.acc-ph], arXiv:1306.6352 [hep-ph], arXiv:1306.6353 [physics.acc-ph], arXiv:1306.6328 [physics.acc-ph], arXiv:1306.6329 [physics.ins-det].
- [14] ILC Higgs report? Snowmass paper.
- [15] D. Dannheim, P. Lebrun, L. Linssen, D. Schulte and S. Stapnes, arXiv:1305.5766 [physics.acc-ph]; H. Abramowicz *et al.* [CLIC Detector and Physics Study Collaboration], arXiv:1307.5288 [hep-ex], Snowmass paper.
- [16] M. Aicheler, *et al.*, CERN-2012-007.
- [17] J-P. Delahaye, C. Ankenbrandt, A. Bogacz, S. Brice, A. Bross, D. Denisov, E. Eichten and P. Huber *et al.*, arXiv:1308.0494 [physics.acc-ph].

- 1325 [18] Y. Alexahin, C. M. Ankenbrandt, D. B. Cline, A. Conway, M. A. Cummings, V. Di Benedetto, E. Eichten
1326 and C. Gatto *et al.*, arXiv:1308.2143 [hep-ph].
- 1327 [19] M. Koratzinos, *et al.*, arXiv:1305.6498 [physics.acc-ph]; M. Koratzinos, A. P. Blondel, R. Aleksan,
1328 P. Janot, F. Zimmermann, J. R. Ellis and M. Zanetti, arXiv:1306.5981 [physics.acc-ph], Snowmass
1329 paper.
- 1330 [20] S. A. Bogacz, J. Ellis, L. Lusito, D. Schulte, T. Takahashi, M. Velasco, M. Zanetti and F. Zimmermann,
1331 arXiv:1208.2827 [physics.acc-ph].
- 1332 [21] W. Chou, G. Mourou, N. Solyak, T. Tajima and M. Velasco, arXiv:1305.5202 [physics.acc-ph], Snowmass
1333 paper.
- 1334 [22] J. L. Abelleira Fernandez, C. Adolphsen, P. Adzic, A. N. Akay, H. Aksakal, J. L. Albacete, B. Allanach
1335 and S. Alekhin *et al.*, arXiv:1211.4831 [hep-ex].
- 1336 [23] S. Dawson, *et al.*, Snowmass paper.
- 1337 [24] H. E. Haber, in *From the Planck Scale to the Weak Scale (SUSY 94)*, P. Nath, T. Taylor, and S.
1338 Pokorski, eds. (World Scientific, 1995) [hep-ph/9501320].
- 1339 [25] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi and M. Spira, Eur. Phys. J. C **71**, 1753 (2011)
1340 [arXiv:1107.5909 [hep-ph]].
- 1341 [26] lattice writeup for Snowmass?
- 1342 [27] CMS Collaboration, Physics Analysis Summary CMS-PAS-HIG-13-005 (2013).
- 1343 [28] V. Barger, L. L. Everett, H. E. Logan and G. Shaughnessy, arXiv:1308.0052 [hep-ph].
- 1344 [29] C. -Y. Chen, arXiv:1308.3487 [hep-ph].
- 1345 [30] E. Brownson, N. Craig, U. Heintz, G. Kukartsev, M. Narain, N. Parashar and J. Stupak, arXiv:1308.6334
1346 [hep-ex], Snowmass paper
- 1347 [31] M. Bardeen, *et al.*, Snowmass paper.
- 1348 [32] A. Kotwal, *et al.*, Snowmass paper.
- 1349 [33] J. Campbell, *et al.*, Snowmass paper.
- 1350 [34] J. Beringer, *et al.* (Particle Data Group), Phys. Rev. **D86**, 010001 (2012).
- 1351 [35] W. -K. Tung, Acta Phys. Polon. B **33**, 2933 (2002) [hep-ph/0206114].
- 1352 [36] C. Anastasiou, L. J. Dixon, K. Melnikov and F. Petriello, Phys. Rev. Lett. **91**, 182002 (2003) [hep-
1353 ph/0306192].
- 1354 [37] M. Czakon, P. Fiedler and A. Mitov, Phys. Rev. Lett. **110**, 252004 (2013) [arXiv:1303.6254 [hep-ph]].
- 1355 [38] R. Aaij *et al.* [LHCb Collaboration], JHEP **1206**, 058 (2012) [arXiv:1204.1620 [hep-ex]].
- 1356 [39] K. Agashe, *et al.*, Snowmass paper.
- 1357 [40] G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori and A. Strumia, JHEP
1358 **1208**, 098 (2012) [arXiv:1205.6497 [hep-ph]].
- 1359 [41] S. Chatrchyan *et al.* [CMS Collaboration], Eur. Phys. J. C **73**, 2494 (2013) [arXiv:1304.5783 [hep-ex]].

- 1360 [42] A. Abdesselam, *et al.*, Eur. Phys. J. C **71**, 1661 (2011) [arXiv:1012.5412 [hep-ph]].
- 1361 [43] Y. Gershtein, *et al.*, Snowmass paper.
- 1362 [44] K. Griest and D. Seckel, Phys. Rev. D **43**, 3191 (1991).
- 1363 [45] ATLAS Collaboration, arXiv:1307.7292 [hep-ex], Snowmass paper.
- 1364 [46] H. Baer, V. Barger, A. Lessa and X. Tata, arXiv:1306.5343 [hep-ph], Snowmass paper.
- 1365 [47] M. Cahill-Rowley, J. L. Hewett, A. Ismail and T. G. Rizzo, arXiv:1307.8444 [hep-ph], Snowmass paper.
- 1366 [48] M. Berggren, F. Brummer, J. List, G. Moortgat-Pick, T. Robens, K. Rolbiecki and H. Sert,
1367 arXiv:1307.3566 [hep-ph].
- 1368 [49] N. Zhou, D. Berge, T. M. P. Tait, L. Wang and D. Whiteson, arXiv:1307.5327 [hep-ex], Snowmass
1369 paper.
- 1370 [50] M. Artuso, *et al.*, Snowmass paper.
- 1371 [51] G. D. Kribs, E. Poppitz and N. Weiner, Phys. Rev. D **78**, 055010 (2008) [arXiv:0712.2039 [hep-ph]].
- 1372 [52] Y. Nomura, M. Papucci and D. Stolarski, Phys. Rev. D **77**, 075006 (2008) [arXiv:0712.2074 [hep-ph]].
- 1373 [53] J. List and B. Vormwald, arXiv:1307.4074 [hep-ex].
- 1374 [54] See, for example, P. Fileviez Perez, T. Han and T. Li, Phys. Rev. D **80**, 073015 (2009) [arXiv:0907.4186
1375 [hep-ph]].
- 1376 [55] D. E. Morrissey and M. J. Ramsey-Musolf, New J. Phys. **14**, 125003 (2012) [arXiv:1206.2942 [hep-ph]].
- 1377 [56] M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D **53**, 2658 (1996) [hep-ph/9507378].
- 1378 [57] N. Arkani-Hamed, A. Delgado and G. F. Giudice, Nucl. Phys. B **741**, 108 (2006) [hep-ph/0601041].